BENHA UNIVERSITY

FACULTY OF VETERINARY MEDICINE

DEPARTMENT OF BACTERIOLOGY, IMMUNOLOGY, AND MYCOLOGY

Written and presented by

Doctor Mohammad Mahmoud Mohammad Yousef Wehaba

Bachelor of veterinary medicine ,kafer el-shiekh university (2003)

Very Good , The Seventh Rank(2012)

Hiring As A Veterinarian Doctor In Animal Health Research Center

Master (Bacteriology, Immunology And Mycology, 2021)

Assistant Researcher Animal Health Institute (2020)

Agricultural Research Center,

Ministry Of Agriculture And Land Reclamation

Arab Republic Of Egypt

Pure microbiology

Branches of microbiology

- **Bacteriology: the study of bacteria.**
- **Mycology: the study of fungi.**
- **Protozoology: the study of protozoa.**
- **Phycology/algology: the study of algae.**
- **Parasitology: the study of parasites.**
- **Immunology: the study of the immune system.**
- **Virology: the study of viruses.**
- **Nematology: the study of nematodes.**

https://en.wikipedia.org/wiki/Branches_of_microbiology

GENERAL BACTERIOLOGY, IMMUNOLOGY, MYCOLOGY

INTRODUCTION TO MICROBIOLOGY

MICROBIOLOGY IS DEFINED AS (THE STUDY OF ORGANISMS TOO SMALL TO BE CLEARLY SEEN BY THE NAKED EYES). THESE ORGANISMS INCLUDE VIRUSES, BACTERIA, ALGEA, FUNGI, ARCHAEA AND PROTOZOA. WHEN STUDYING MICROORGANISMS , MICROBIOLOGISTS ARE CONCERNED WITH CHARACTERISTICS AND FUNCTIONS SUCH AS MORPHOLOGY ,CYTOLOGY, PHYSIOLOGY , ECOLOGY , TAXONOMY , GENETICS , AND MOLECULAR BIOLOGY .

علم البيئة ECOLOGY التصنيف الحيوى TAXONOMY

Leeuwenhoek

(Live 1632 -1723)

Is The First Person To Use The

Microscopes To Observe Microbes:

He Made Small Handheld Microscopes:

((He Called Microorganisms

As Animalcules))

خلايا بدائيات النوى خلية بدائية النواة Procaryotes

TAXONOMY AND CLASSIFICATION OF PROCARYOTES

HEAKEL (1866) WAS THE FIRST TO CREATE ANATURAL KINGDOM FOR MICROORGANISMS, HE PLACED ALL UNICELLULAR (MICROSCOPIC) IN ANEW KINGDOM CALLED PROTISTA, SEPERATED FROM PLANTS (PLANTAE) AND ANIMALS (ANIMALIA) , WHICH WERE MULTICELLULAR (MACROSCOPIC) ORGANISMS .

PROTISTA : UNICELLAR MICROSCOPIC ORGANISMS

PLANTAE : MULTICELLULAR MACROSCOPIC ORGANISMS

ANIMALIA : MULTICELLULAR MACROSCOPIC ORGANISMS

IN (1950) :

THE DEVELOPMENT OF ELECTRON MICROSCOPE

THE ELECTRON MICROSCOPE MORE ONE MILLION POWERFUL THAN THE LIGHT MICROSCOPE.

SO BY THE ELECTRON MICROSCOPE THEY FIND ANOTHER KINGDOM CALLED MONERA (MONERES) : INCLUDES MICROSCOPIC UNICELLULAR ORGANISMS BUT NOT HAVE THE NUCLEAR MEMBRANE .

PROTISTA : REMAINED AS AKINGDOM OF UNICELLULAR EUCARYOTIC MICROORGANISMS .

WHITTAKER : REFINED THE ECOSYSTEM INTO FIVE KINGDOMS IN (1967) BY IDETIFYING THE FUNGI AS ASEPERATE MULTICELLULAR EUOCARYOTIC KINGDOM OF ORGANISMS , DISTINGUISHED BY THEIR ABSORPTIVE MODE OF HETEROTROPHIC MODE OF NUTRITION .

يتكون النظام البيئى من خمس مملكات حيوية :

THE FIVE KINGDOMS OF THE ECOSYSTEM:

1-PROTISTA : UNICELLAR MICROSCOPIC ORGANISMS

2-PLANTAE : MULTICELLULAR MACROSCOPIC ORGANISMS

3-ANIMALIA : MULTICELLULAR MACROSCOPIC ORGANISMS

4- MONERA : UNICELLULAR MICRORGANISMS NOT HAVE NUCLEAR MEMBRANE.

5- FUNGI : MULTICELLULAR OF EUCRAYOTIC ORGANISMS ITS LENGHTH 4 MICROMETERS IN LENGTH .

BACTERIA IS 1 MICROMETERS IN LENGTH

FUNGI IS 4 MICROMETERS IN LENGTH

FIVE KINGDOMS OF THE ECOSYSTEM AS FELLOW :

1. **ANIMALIA :INCLUDES VERTEBRATES AND ARTHROPODES**

2. **PLANTEA : INCLUDES GREEN ALGEA**

3. **FUNGI : INCLUDES MOLDS AND YEASTS**

4. **MONERA : INCLUDES ARCHAEA ,BACTERIA, MYCOPLASMA**

5. **PROTESTA : INCLUDES RED ALGEA , AMOEBAS, WATER MOLDS .**

What are 5 characteristics of fungi?

Characteristics of Fungi

- Fungi are eukaryotic, non-vascular, non-motile and heterotrophic organisms.
- They may be unicellular or filamentous.
- They reproduce by means of spores.
- Fungi exhibit the phenomenon of alternation
- of generation.(Absorptive mode of Nutrition)
- Fungi lack chlorophyll and hence cannot perform photosynthesis.
- Have two types of reproduction sexual and asexual type

- **A Reportable Animal Diseases Are That, By Law, Must Be Reported To State Animal Health Officials.**

المرض المشتبة به	نوع العينة المراد العزل منها	المادة الحافظة	طريقة النقل المناسبة ودرجة الحرارة المطلوبة	المدة المسموح بها حتى الوصول إلى معهد الدقى
1-الحمى القلاعية (FMD)	حويصلة كاملة- بثرات- سائل الحويصلة	Glycerin buffer	فى ثلاجة نقل العينات محتوية على أكياس ثلج خاصة بحيث تكون(4C) كل عينة فى كيس منفصل	24 ساعة
2-حمى الوادى المتصدع(RFV)	دم على مانع تجلط – أجنة مجهضة (كبد- طحال-غدة ليمفاوية)		كما سبق	24 ساعة
3-الجلد العقدى (LSD)	عقدة جلدية – عقدة ليمفاوية – طحال	Transport buffer	كما سبق	24 ساعة
4-طاعون المجترات الصغيرة (PPR)	دم على مانع تجلط أثناء إرتفاع درجة الحرارة –بثرات من الفم – مسحات من العين والأنف والفم –أحشاء (الرئة غدد ليمفاوية – الأمعاء الطحال)	EDTA Heparin	كما سبق	4-6ساعات
5-التسمم الدموى (Hemorrhagic septicemia) (Pasteurella multocida)	مسحات أنفية – أحشاء داخلية – دم –سيرم		كماسبق	4-6ساعات
6-حمى الثلاث أيام	دم على مانع تجلط – غدة ليمفاوية - طحال	Preservative media PBS	كما سبق	24 ساعة
7- الإلتهاب الشعبى البقرى (IBR)	مسحات أنفية-غدة ليمفاوية-جزء من الرئة –جزء من القصبة الهوائية	Preservative media PBS	كما سبق	24ساعة
8-الإسهال المعدى (BVD)	مسحات أنفية- مسحات شرجية- سيرم-غدة ليمفاوية – أحشاءداخلية	Preservative media PBS	كما سبق	24ساعة

9-أمراض العجول الرضيعة	عينات براز-سيرم-أحشاءداخلية-مسحات أنفية ودم	Preservative media PBS	كما سبق	24ساعة
10-التسمم المعوى (Enterotoxaemia)(clostridium perfringes type D)	الأمعاءالدقيقة (مربوطة ولا يقل طولها عن 15 سم) –فص كامل من الكبد به الإصابة – كلية كاملة		كما سبق	4-6ساعات
11-التفحم العضلى والأوديما الخبيثة Black Leg (black quarter) Malignant Edema (clostridium chauvoei)	العضلة المصابة المتفحمة –سائل إرتشاحى من مكان الإصابة		كما سبق	4-6ساعات
12-المرض الأسود(Black Disease) (clostridium chauvoei)	فص كامل من الكبد به إصابة –الأمعاء الدقيقة		كما سبق	4-6ساعات
13-باسيلارى هيموجلوبين يوريا (Bacillary Hemoglobinuria (Red urine)(clostridium novyai)	فص كامل من الكبد به الإصابة		كما سبق	4-6ساعات
14-تعفن الحافر (fusobacterium necrophorum) (Foot Rot)	مسحات من الحافر المصاب بعد تنظيفه جيدا من الأنسجة الميتة	المسحات على ناقل (Cary-Blair Medium)	كما سبق	4-6ساعات
15- التسمم الغذائى Unbalanced Ration,Toxic Chemical Addetives,Mycotic Infections And Mycotoxins	-الأعلاف (لا تقل العينة عن 50-100جرام) -السيلاج (لا تقل العينة عن 50-100 جرام)- المياة (لا تقل عن 200 مل)		كما سبق	لا تزيد عن 12 ساعة بطريقة الحفظ المذكورة

الأمراض الوبائية المتوطنةوالكشف عنها ووجب تحصينهاداخل القطر المصرى .

البروسيلا تعتبر من أهم أمراض طوارئ الإتصال وقد تم نسيانها والإسبريرجلوسس تعتبر من أهم الأمراض الفطرية وتسبب نكسات مزرعية تعتبر من أهم أمراض طوارئ الإتصال وقد تم نسيانها

PBS Phosphate Buffer Saline A Reportable Animal Disease Is One That, By Law, Must Be Reported To State Animal Health Officials.

أمراض العجول الرضيعة تشمل :

Suckling Calves Fatal Diseases

Corona,Parainfleunza,Pneumoenteritis Viruses May Be Mixed
E.Coli,Salmonella,Mycoplasma Fatal Infections.

العينات يجب أن تكون طازجة من الحيوانات النافقة بعد نفوقها مباشرة وإذا لوحظ أى تغيرات رمية فى الحالة أوالحالات المعتبرة بالكشف لا يؤخذ منها أى عينات للحكم أو التعويض لتأكيد إدعاؤه وطلبه وإذا بلغ بوجود تسمم كيماوى يتم إحالة الملف إلى الطب الشرعى للبت فيها ولا يوجد أى إشكال لو كانت الحالات على قيد الحياة وتفضل فى أخذ العينات والحالات النافقة فقط لتأكيد المسببات المرضية الوبائية المتوطنة ويجب أن يقدم الطبيب البيطرى الخاص بالمزرعة ما يدل على التحصين الفعلى الحكومى ويتم التأكيد الحكومى الإدارى والفحص المناعى .

ملحوظة هامة : يجب الإفادة عن الموقف التحصينى لتلك الأمراض عند إرسال أى عينات للفحص

تحريرا فى : / /

يعتمد

أ.د وكيل المعهد للتشخيص المعملى وصحة الأغذية

Some of Important bacterial diseases

Streptococcus	**pneumonia , pharyngitis , cellulitis**
Staphylococcus	**abscess of skin and other organs**
Bacillus anthraces	**anthrax**
Clostridium	**tetanus , gas gangrene , botulism**
Coryne bacterium	**diphtheria**
Listeria	**meningitis**
Actinomyces	**actinomycosis**
Nocardia	**nocardiosis**
Neisseria	**gonorrhea , meningitis**
Haemophilus	**meningitis**
Bortedella	**whooping cough**
Legionella	**pneumonia**
Brucella	**brucellosis**
Francisella	**tularemia**
Pasteurella	**cellulitis**
Yersinia	**plaque**
Escherichia coli	**urinary tract infection , diarrhea**

Enterobacter	**urinary tract infection**
Serratia	**pneumonia**
Klebsiella	**urinary tract infection**
Salmonella	**enterocolitis , typhoid fever**
Shigella	**enterocolitis**
Proteus	**urinary tract infection**
Campylobacter	**enterocolitis**
Helicobacter	**gastritis , peptic ulcer**
Vibrio	**cholera**
Pseudomonas	**urinary tract infection , pneumonia**
Bacteroids	**peritonitis**
Mycobacterium	**tuberculosis , leprosy**

Rickettssia (rocky mountain spotted fever , typhus , Q fever)

Chlamydia	**urethritis , trachoma , psittacosis**
Treponema	**syphilis**
Borrelia	**lyme disease**
Leptospira	**leptospirosis**
Mycoplasma	**pneumonia**

الأهمية الطبية لمادة البكتيريا والمناعة والفطريات فى الحياة العامة

MEDICAL IMPORTANCE OF BACTERIOLOGY AND IMMUNOLOGY AND MYCOLOGY IN OUR LIFE

MYCOTIC DISEASES ARE CAUSING SEVER BIOLOGICAL SYSTMIC CELLULAR DESTRUCTIONS AND DEATHS IN LIVING ANIMALS AND BIRDS.

IMMUNOLOGY MUST ESTABLISH TO CONFIRM THE HEALTH DEGREE AND READINESS TO VACCINATIONS AND IMMUNIZATIONS IN HEALTHY AND DISEASED ANIMALS AND BIRDS.

Classification of medically important bacteria

1-RIGID, THICK- WALLED CELLS

2- FLEXIBL, THIN WALLED CELLS

3- WALL LESS – CELLS

1- RIGID, THICK- WALLED CELLS BACTERIA :

INCLUDES:

A- FREE- LIVING BACTERIA (EXTERA CELLULAR BACTERIA)

B- NON-FREE-LIVING BACTERIA (OBLIGATE INTERACELLAR BACTERIA)

(As Rickettsia And Chlamydia)

A- FREE- LIVING BACTERIA (EXTERA CELLULAR BACTERIA)

DEVIDED ACCORDING TO GRAM STAIN INTO:

1- Gram Positive Bacteria (Stained Appear Blue Colour)

2- Gram Negative Bacteria (Not Stained Just Avery Little Appear Red Colour)

3-Acid Fast Bacteria E.G Tubercle Bacilli

1- Gram Positive Bacteria : Have Two Shapes : Cocci And Bacilli

2- Gram Negative Bacteria : Have Two Shapes : Cocci And Bacilli

1- Gram Positive Bacteria :

A- Cocci B- Bacilli

A- Cocci Likes Staph And Strept

B- Bacilli Include : Spore Forming And Non Spore Forming

1- Spore Forming Include :

a- Aerobic Bacteria Like Bacillus Anthraces

b- Anaerobic Bacteria Like Clostridia

2- Non Spore Forming Divided Into:

a- Non Filamentous Like Coryn Bacteria, Listeria

b- B-Filamentous like Actinomyces

2- Gram Negative Bacteria : Have Two Shape Cocci And Bacilli

A- Cocci Like Nisseria

B- Bacilli Include :

a- Aerobic (Psudomonas)

b- Facultative Anaerobic (Straight – Curved)

c- Anaerobic Bacteria (Bacteroides)

Facultative Anaerobic Bacteria : (Straight – Curved)

a- Straight (Respiratory Bacteria- Zoonotic Bacteria-Enteric Bacteria)

Respiratory Likes Haemophilus

Zoonotic Likes Brucella , Pasteruella , Yersenia

Enteric Likes E.Coli , Salmonella, Shigella ,Klebsiella, Proteus

b- Curved Likes Campylobacter , Vibriocholera, Hilicobacter

2- Flexible , Thin Walled, Cells Likes Leptospira

3- Wall-Less Cell Wall Likes Mycoplasma

Classification Of Important Bacterial Diseases:

Rigid,Thick -Walled Cells	Flexible,Thin-Walled Cells (Spirochetes) :As 1-Treponema Causing Syphilis 2-Borrelia Causing Lyme Disease 3- Leptospira Causing Leptospirosis	Wall-Less Cells (Mycoplasma) Causing Pneumonia

Rigid thick walled cells	
Extera-cellular free living bacteria	Obligatenteracellular Non-free-living bacteria : as

	1-Rickettssia Causing **(Rocky Mountain Spotted Fever , Typhus , Q Fever)** **2-Chlamydia Causing** **(Urethritis , Trachoma , Psittacosis)**

Free Living Bacteria (Extra And Intra Cellular Parasite)		
1-Gram Positive **Blue In Color**	**2-Gram Negative** **Red Or Pink Color**	**3-Acid Fast As:** **Mycobacterium Causing :** **Tuberculosis ,** **Leprosy**

Gram positive bacteria

Cocci (spherical shape)	**Bacilli (rods)(cigar shape)**	
Streptococcus causing pneumonia , pharyngitis , cellulitis **Staphylococcus abscess of skin and other organs**	**sporeforming**	**non-sporeforming**
	1- Aerobic as Bacillus anthraces causing anthrax **2-anaerobic as Clostridium causing tetanus , gas**	**1-Non Filamentous As** **A- Coryne Bacterium Causing Diphtheria** **B-Listeria Causing**

	gangrene , **botulism**	**Meningitis** **2- Filamentous As Actinomyces Causing Actinomycosis**

Gram Negative Bacteria	
Cocci (spherical shape)As Neisseria Causing Gonorrhea , Meningitis	**Bacilli (Rods)(Cigar Shape) Includes** **1-Aerobic** **2-Facultativeanaerobic** **3-Anaerobic**

Gram Negative Bacilli (Rods)(Cigar Shape)		
Aerobic as **Pseudomonas causing : urinary tract infection , pneumonia**	**Facultative Anaerobic May Straight Or Curved.**	**Anaerobic as** **Bacteroids causing : peritonitis**

Curved Facultative Anaerobic Gram Negative Bacilli (Rods) (Cigar Shape)		
Campylobacter	**Causing**	**Enterocolitis**
Helicobacter	**Causing**	**Gastritis , Peptic Ulcer**
Vibrio	**Causing**	**Cholera**

Straight Facultative Anaerobic Gram Negative Bacilli (Rods) (Cigar Shape)		
Respiratory **Haemophilus meningitis** **Bortedella whooping cough** **Legionella pneumonia**	**zoonotic** **Brucella brucellosis** **Francisella tularemia** **Pasteurella cellulitis** **Yersinia plaque**	**Enteric and related** **Escherichia coli urinary tract infection , diarrhea** **Enterobacter urinary tract infection** **Serratia pneumonia** **Klebsiella urinary tract infection** **Salmonella enterocolitis , typhoid fever** **Shigella enterocolitis** **Proteus urinary tract infection**

Written And Presented By

Doctor Mohammad Mahmoud Mohammad Youssef Wehaba

FACULTATATIVE ANAEROBIC GRAM NEGATIVE RODS

FAMILY: ENTEROBACTERIACEAE

THE DEFINITION OF ENTEROBACTERIACEA:

Intestinal Bacteria Describe Large Biochemically Antegenically Related Groups Of Organisms

1-Some Of Them Are Pathogenic Potential To Man, Animals, Insects And Plants.

2- It Contains 4 Tribes, 12 Genera On The Basis Of DNA Relatedness.

3-Generally The Members Of Family Enterobacteriaceae Is Composed Of Gram Negative Rods Which May Be Motile By Peritrichous Flagella Or Non-Motile.

4- They Are Non -Spore Forming And Not Acid Fast.

5- Most Of Them Are Non Capsulated But Some Have Capsules Or Microcapsules (Slime Layer).

6- Enterobacteriaceae Members Grow On The Presence And The Absence Of Free Oxygen Because Facultative Anaerobes Have The Presence Of Respiratory And Fermentative Metabolisms.

5- They Grow Well On Ordinary Media.

6- They Are Similar In The Morphological And Cultural Characters.

7-Moreover All Species Contain The Enterobacterial Common Antigen.

8- Biochemically They Produce Acid And Gas Due To The Fermentation Of Glucose Or Different Types Of Carbohydrates.

9- All Species Are Catalase Positive .

10-Reduce Nitrate To Nitrite.

11- Oxidase Positive .

12- According To Their Ability To Ferment Lactose They Were Differentiated Into Biochemically Into Two Divisions :

1 - Lactose Fermenting Enterobacterial Genera.

2-Non Lactose Fermenting Enterobacterial Genera .

This Classification Depends On The Cultivation Of The Species Of Family Enterobacteriaceae On One Of The Following Differential Selective Media :

1-MacConkey agar (lactose-bile neutral red agar)

2-S.S media (salmonella- shigella agar)

3- Desoxycholate cirate agar

4- Hegkton media

Different genera enterobacteriaceae according to their ability to ferment lactose :

Lactose fermenter :

Genus	Habitate
E.coli	Intestine
Enteobacter	Intestine , water, Intestine
Citrobacter	Water , soil, Intestine
Yersenia	Water , Intestine
Serratia	Water , soil, Intestine

Non- lactose fermenter :

Genus	Habitate
Shigella	Intestine ,water, soil
Salmonella	Intestine , water , soil
Hafnia	Intestine, water
Proteus	Intestine, water , soil
Morganella	As Proteus

Genus: Eshcerichia

Species: Escerichia Coli (E.Coli)

Synonym: Bacterium Coli (E.Coli)

Definition: Escherichia Coli Is A Normal Inhabitant Of The Lower Part Of Intestine Of All Warm Blooded Animals. It Is Rarely Present In Stomach And In Few Numbers In The Anterior Part Of Intestine. Carnivores And Omnivores Usually Harbor E.Coli In Greater Number Than Herbivore

Under Certain Conditions Of Stress May Undergo To Be Potential Pathogenic Causing Sever Diseased Conditions.

Specific Diseases Caused By Enteropathogenic E.Coli In Different Animal Species:

Species	Disease	Symptoms
Childrens	Coli Septicemia	Diarrhea , Death
Newly Born Animals	Colibacillosis, White Or Calf Scour	Septicemia And Rapid Death

Microscopic Characters :

E.Coli Is Coccobacilli To Long Rods , Arranged Singly Or In Short Chains Especially In Mucoid Colonies Where It Had A Slime Layer Of Polysaccharides .

It Is Motile And Non – Sporulated .

Cultural Characters :

(Nutrient Broth-Nutrient Agar-Blood Agar-Macconkey Agar)

-In Nutrient Broth Growth Is Rapid , Gives Regular Turbidity Within 12-18 At 37 C With Heavy Powdery Sediment After 24 Hours .

-On Nutrient Agar Growth May Be Or Together

1-Smooth Surface Colonies Occurs After 24 Hours,

-The Colonies Are Grayish In Color , Circular , Convex 2-3mm In Diameter .

2- Mucoid Colonies Appear Due To The Presence Of Slime Layer And Larger In Size Than Smooth One .

-On Blood Agar The Surface Colonies Are Larger Some Are Hemolytic And Others Are Non-Hemolytic

-On Macconkey Agar It Gives Round Pink Colonies, Medium Sized Opaque Colonies

I.E Lactose Fermenter

Environmental Characters :

(Viability And Resistance) :

-It May Survive For Weeks Or Months It Contaminated Water (Sewage Pollution) , Dwellings In The Animal Houses In The Presence Of Mucous Or Feces . Some Strains Resist Dryness.

-It Is Susceptible To Physical And Chemical Agents. E.Coli Is Killed By Heating At (55-60 C) For 30 Minutes, Within 10 Minutes By Autoclaving.

Antigenicity Of Escherichia Coli :

1-Somatic Antigens (146 O- Antigens) 2-Capsular Antigens (91 K –Antigens) 3- Flagellar Antigens (49 H- Antigens) :

1-Somatic Antigens : (146 O- Antigens) :

E.Coli Strains Have 146 O-Antigens In Their Bodies . Each Strain Carries Onely One Type Of O-Antigen . They Are Composed Chemically Of Polysaccharides – Phospholipids –Protein Complex. The O –Antigens Are

Designated By Numbers As O1, O2, O3,O4 ,O5 – O 145 Etc .

The O – Antigens Are Heat Stables.

2- Capsular Antigens (91 K –Antigens) :

These 91 K-Antigens Occur As A Slime Layer Or Microcapsule On Most Enteropathogenic E.Coli Strains. Chemically They Are Composed Of Poly Saccharides . K-Antigens Prevent Agglutination Between O-Antigens And Their Specific Homologous Antisera.

The K – Antigens Are Differentiated Into

3 Types According To Thermo Stability As Follows :

L-Antigen : Heat Labile

A- Heat Stable

B- Antigen : Lipopoly Saccharide Destroyed By Autoclaving

The K-Antigen Is Very Important In Serotyping Of E.Coli As Well As In The Mechanism Of Pathogenesis As K 88 And K 99 .

These K 88 And K 99 Mediate Adhesion Of The Organism To The Microvilli Of The Intestinal Epithelial Cells And Enhance Colonization .

3-Flagellar Antigen (49 H-Antigens) :

There Are 49 H-Antigens Which Is Composed Of Protein . Frequently It Is Poorly Developed .

It Has Weak Role In Serotyping Of E.Coli.

Serotyping Identification : -

For Serotyping Of E.Coli , Identification Of O And K Antigens Is Applied By Slide Agglutination Test To Determine Its Antigenic Structures As O20/B6 , O55/B5 , O88/B4 , O111/B4 .

(Ent.Path)Toxins Of E.Coli (E.Coli Products) :

1-Enterotoxin 2- Colicins

1-Enterotoxin:

This Substances Is Produced By Entero Pathogenic Strains Of E. Coli , And Causes Diarrhea In Man And Animals . Injection Of E .Coli Enterotoxins Into The Ligated Intestinal Loops Of Rabbits Induced Swelling , Oedema And Accumulation Of Exudate In The Injected Loop. The Enterotoxins Are Not Produced By All E.Coli Strains But Only The Enteropathogenic.

E.Coli Enterotoxin Is Composed Of Two Fractions:

A- Heat Stable Fraction.

B-Heat Labile Fraction.

A-Heat Stable Fraction : It Is Exteracellular In The Culture . It Is Resistant To Heating For 60 C But Destroyed By Autoclaving .

B- Heat Labile Fraction: It Is Inactivating By Heating At 60 C For 30 Minutes It Is Antigenic And Occurred Within Bacterial Cells . It Could Be Neutralized By Specific Antisera .

2- Colicins :

It Is E.Coli Products And Composed Chemically Of Protien . It Has Bactericidal Activitiy . Colicins Productions Is Very Important In Differentiation Of E.Coli Strains Into :

Colicins Positive E.Coli.

Colicins Negative E.Coli.

Laboratory Diagnosis:

- The Isolation Of E.Coli May Be No Value As E.Coli Are Normal Inhabitant Of The Intestinal Tract, Unless Other Seven Factors Are Considered Including:

1-Detailed History Of The Case, Symptoms And Lesions With Differential Bacteriological Diagnosis On The Other Causative Agents Of Digestive Disorders

2-Purity Of The Growth Of E.Coli On The Primary Cultures Note The Growth Of Lactose Fermenting Colonies On Macconkey Agar .

3-Site Of Isolation From Various Organs Especially From The Parenchymatous Organs As Liver , Spleen , Mesenteric Lymph Nodes And Lung Tissues Indicating Colibacillosis.

4- Biochemically Identification Of Isolated Suspected E.Coli The Suspected E.Coli Colonies As Lactose Fermenter On Macconkey Media Can Be Tested Biochemically

And Must Be Differentiated From The Other Lactose – Fermenter Species By The Following Reactions (Indol , Methyle Red , Vp , Citrate , Urea Hydrolysis , H2s, Motility , Kcn) (Voges Proskauer)(Potassium Cyanide)

E.Coli Are Indol And Methyle Red Positive Only

5-Serotyping Of E.Coli If The Isolating Colonies, Are Biochemically Typical Of E.Coli, Serotyping Must Be Performed Using Polyvalent O, K Anti-Sera To Determine The Antigenic Structure Of The Isolated Strain Of E.Coli.

6-Final Diagnosis Depends Mainly Upon The Isolation Of E.Coli From Various Fresh Organs In Case Of Septicemia Of The Same Antigenic Structure.

7-It Is Very Essential That The Same Serotype Of E.Coli Must Be Isolated From Feces Or Intestinal Content As Well As The Internal Organs Of The Same Case .

So There Two Points Of View Of Doctor Mohammad Mahmoud Mohammad Youssef Wehaba :

1- (Net Source E.Coli More Pathogenic Than Shigella And Klebsiella Because The Both Are Not Motile But Escherichia Coli Have Very Strong Peritrichous Flagellae So This Is Genus Had Vital Accommodated Advanced Genetically Assortment).

2- (From My Study 2001 I Am Suring On Pathological Basis There Is A Difference Between Commensals And Pathotypes So We Must Depend On Genotyping) (You Ask How)(Ok In The Stress The Inter Cellular Spaces And Pores Will Increases So Limits ,Borders Are Open So Commensals Without Any Efforts Invading The Cells Replicates And Colonized Due To Leaked Nutrients And Weak Innate Immunity)

(Commensals Only Causing Travelling Diarrhea)

(Pathotypes Causing Adiseases Whatever The Health State)

BENHA UNIVERSITY
FACULTY OF VETERINARY MEDICINE
DEPARTMENT OF CLINICAL PATHOLOGY
Written and presented by
Doctor Mohammad Mahmoud Mohammad Yousef Wehaba
Bachelor of veterinary medicine ,kafer el-shiekh university (2003)
Very Good , The Seventh Rank(2012)
Hiring As A Veterinarian Doctor In Animal Health Research institute
Master in (Bacteriology, Immunology And Mycology, 2021)
Assistant Researcher Animal Health Institute (2020)
Researcher assistant animal health research institute (2022)
Agricultural Research Center,
Ministry Of Agriculture And Land Reclamation
Arab Republic Of Egypt

BODY FLUIDS EXAMINATION

ELECTROLYTES AND TRACE ELEMENTS :

1- WATER METABOLISM
2- CYSTATIN C

1-WATER METABOLISM

1-WATER CONSTITUTES ABOUT 45-70 % OF TOTAL BODY WEIGHT .

2- BODY FLUIDS ARE DIVIDED IN TWO COMPARTMENTS:

(A)- INTRA CELLULAR (ICF), WHICH WHICH CONSTITUTES 65-75 % OF TOTAL BODY WATER AND 35-45 % TOTAL BODY WEIGHT .

(B)- EXTERA CELLULAR FLUIDS (ECF) CONSTITUTES ABOUT 25 % OF TOTAL BODY WATER AND 15 % OF TOTAL BODY WEIGHT .

3-)- EXTERA CELLULAR FLUIDS (ECF) FOUND IN 3 LOCATIONS :

(A)- INTRA VASCULAR (PLASMA)

(B)-INTRASTITIAL (LYMPH)

(C) – TRANS CELLULAR (IT INCLUDES : CSF ,JOINT FLUIDS ,PERITONEAL PERICARDIAL, PLEURAL AND LIQUID IN INTESTINE).

4-DECREASED TOTAL BODY WATER LEADS TO DEHYDERATION

5-DEHYDERATION IS MANIFESTED BY :

(A)-DECRASED BODY WIEGHT

(B)LOSS OF SKIN ELASTICITY

(C)-DRYNESS OF MUCOUS MEMBRANE

(D)-RETRACTIONOF EYES OF THE EYES

6- ELECTROLYTES AND WATER MOVES FREELY AND CHARGELY BETWEEN THE ICF AND ECF .

MOVEMENT OF WATER IS DEPENDANT UPON ON HYDROSTATIC PRESSURE AND OSMOTIC PRESUURE OF ECF PROTIENS

7- OSMOTIC PRESSURE OF ANY FLUIDS IS DIRECTLY RELATED TO THE NUMBER OF PARTICLES IN A UNIT OF SOLVENT AND IS MEASURED BY OSMOLE OR MILLIOSMOLE .

8- OSMOLE IS THE MOLECULAR WEIGHT OF THE SUBSTANCE MULTIPLIED BY THE NUMBER OF IONS INTO WHICH IT DISSOCIATES .

9- THE NORMAL OSMOLALITY OF ECF IS ABOUT 300 MOSM /KG

10-ECF CONTAINS SODIUM NA , CHLORIDE CL AND BICARBONATE HCO3

11- ICF CONTAINS K POTTASIUM ,MG MAGNESIUM ,ORGANIC PHOSPHATES AND PROTIEN.

12-SERUM OSMOLALITY CAN BE ESTIMATED BY USING THE CONCETRATION OF NA SODIUM , AND K POTTASIUM AS FOLLOW :

MOSM / KG = 2(NA +K) +BUN/2.8+GLUCOSE/18

13-SERUM OSMOLALITY IS USED AS AMEASURE OF ECF OSMOLALITY .

14-HYPER OSMOLALITY IS FOUND IN CASE OF :

(A)-HYPERNATREMIA

(B)-ACCUMLATION OF ENDOGENOUS SOLUTES SUCH AS GLUCOSE ,KETONE BODIES , AND UREA

15- HYPOOSMOLALITY IS FOUND IN CASE OF HYPONATERMIA

16-OSMORECEPTORS SENSE CHANGES IN OSMOLALITY . WITH HYPER OSMOLALITY (HYPONATREMIA) , OSMORECEPTORS STIMULATES VASOPRESSIN OR ADH SECRETION FROM THE PITUITARY GLAND AND STIMULATE THIRST . THIRST IS STIMULATE BY AS ALITTLE AS

1-2 % CHANGES IN OSMOLALITY .THE END RESULTS IS WATER RETENTION BY THE KIDNEY AND INCREASED WATER INTAKE . INCREASES IN FREE WATER WILL THUS WILL REDUCE SODIUM CONCENTERATION ,OPPOSITECHANGES OCCUR WITH HYPOOSMOLALITY.

17-BARORECEPTORS ARE SENSITIVE TO CHANGES IN EFFECTIVE CIRCULATING VOLUME (ECV) THE ECV IS THAT PART OF THE EXTRACELLULAR FLUID THAT IS IN THE ARTERIAL SYSTEM AND IS EFFECTIVELY PERFUSING THE TISSUES . IT USUALLY VARIES DIRECTLY WITH ECF VOLUME . WITH HYPOVOLEMIA (DECREASED ECV), BAROECEPTORS STIMULATES THE RENIN-ANGIOTENSIN SYSTEM , THE END RESULT BEING MINERALOCORTICOID (ALDOSTERONE) RELEASE

FROM THE ADRENAL CORTEX. ALDOSTERONE INCREASED ABSORPTION OF NACL AND PROMOTES THE EXCRETION OF POTASSIUM AND HYDROGEN IN THE DISTAL TUBULES OF THE NEPHRON , NACL RETENTION PROMOTES WATER REABSORPTION , THIS CORRECTING THE HYPOVOLEMIA . HYPOVOLEMIA ALSO STIMULATES THIRST (A DECREASE IN ECV OF 7-10 % IS REQUIRED FOR STIMULATION).

OPPOSITE CHANGES OCCUR WITH HYPER VOLEMIA

ANTIDIURETIC HORMONE

ADH ACTS ON CHIEF CELLS IN COLLECTING TUBULES , INCREASING PERMEABILITY OF THE TUBULES TO WATER AND UREA . ADH ACTS VIA ADENYLE CYCLASE WHICH OPEN CHANNELS IN THE TUBULES ,CALLED AQUA-PORES , FACILITATING WATER ABSORPTION .ADH IS SECRETED BY HYPOTHALAMUS IN RESPONSE TO OSMOTIC (MAINLY DETERMINED BY NA SODIUM) AND NON OSMOTIC (VOLUME DEPLETION ,NAUSIA ,FEAR,ANXIETY,PAIN,EXERCISE AND DRUGS) STIMULI.

SEVERAL THINGS INHIBIT THE ACTION OF ADH ON RENAL TUBULES AND THUS AFFECT CONCENTRATING ABILITY .

THESE INHIBITORS OF ADH INCLUDE :

(1)-HYPOKALEMIA

(2)-HYPERCALCEMIA

(3)- CORTICOSTEROIDS

(4-)-PROSTAGLANDIN E

IN ANY OF THE ABOVE CONDITIONS URINE CONCENTRATING ABILITY IS IMPAIRED FOR REASONS OTHER THAN PHYSIACL LOSS OF FUNCTIONAL NEPHRONS .

KIDNEYS

CYSTATIN C

SPECIMEN: SERUM OR PLASMA (HEPARIN)

METHODOLOGY: IMMUNOASSAY

REFERENCE VALUE: 0, 53 - 0, 95 mG /L

Description : CYSTATIN C IS ACYSTIENE PROTIENASE INHIBITOR , ASMALL MOLECULE THAT IS PRODUCED BY NUCLEATED CELLS . SERUM CYSTATIN C IS FILTERED OUT OF BLOOD BY THE KIDNEYS AND THAT SERVES AS AMEASURE OF KIDNEYS FUNCTION. ITS MOLECULAR MASS ALLOWS IT TO BE FREELY FILTERED BY THE GLOMERULAR MEMBRANE IN THE KIDNEYS. SINCE IT IS FORMED AT ACOSTANT RATE AND FREELY FILTERED BY THE HEALTHY KIDNEYS, CYSTATIN C IS AGOOD MARKER OF RENAL FUNCTION. SERUM CONCENTERATIONS OF CYSTATIN C ARE ALMOST TOTALLY DEPENDANT ON GLOMERULAR FILTERATION (GFR) .AREDUCTION IN GFR CAUSES ARISE IN THE CONCENTERATION OF CYSTATIN C. CYSTATIN C IS ABETTER MARKER OF THE GLOMERULAR FILTERATION RATE AND HENCE OF KIDNEY FUNCTION THAN CREATININE WHICH WAS THE MOST COMMONLY USED MEASURE OF KIDNEY FUNCTION. THE LEVELS OF CYSTATIN C ARE INDEPENDENT OF WEIGHT AND HEIGHT ,MUSCLE MASS,AGE, (OVER AYEAR OF AGE) ,SEX,AND FACTORS WHICH HAVE BEEN DOMENSTERATED TO AFFECT CREATININE VALUES,IN ADDITION,ARISE IN CREATININE DOES NOT BECOME EVIDENT UNTIL THE GFR HAS FALLEN BY APPROXIMATELY 50%. MEASUTEMENTS CAN BE MADE AND INTERPRETED FROM ASINGLE RANDOM SAMPLE. CYSTATIN C MAY IS USED AS AN ALTERNATIVE TO CREATININE AND CREATININE CLEARANCE TO SCREEN FOR AND MONITOR KIDNEY DYSFUNCTION IN THOSE WITH KNOWN OR SUSPECTED KIDNEY DISEASES. IT MAY BE ESPECIALLY USEFUL IN THOSE CASES WHERE CREATININE MEASUREMENT IS NOT

APPROPRIATE: FOR INSTANCE IN THOSE WHO HAVE LIVER CIRRHOSIS, ARE VERY OBESE, ARE MALNOURISHED OR HAVE AREDUCED MUSCLE MASS. MEASURING CYSTATIN C MAY ALSO IS USEFUL IN THE EARLY DETECTION OF KIDNEY DISEASE WHEN OTHER PARAMETERS MIGHT STILL BE NORMAL, ESPECIALLY IN THE ELDERLY.

كلية معتمدة 2013

Benha University
Faculty of Veterinary Medicine
Bacteriology, Immunology, Mycology Department

Occurrence of foodborne pathogen in some types of soft cheese in local market

A Thesis presented by
Faculty of Veterinary Medicine
Benaha University

Presented By

Mohammed Mahmoud Mohammed Yousef Wehaba
(BVSc., Kefier Elsheikh University, 2003)
(Bacteriology, Immunology, Mycology)

Under Supervision of

Prof. Dr. Ashraf Awad Abd El tawab
Professor of Bacteriology, Immunology, Mycology Department
Faculty of Veterinary Medicine
Benha University

Prof. Dr. Amal Mostafa Eid
Chief Researchers
Animal Health Research Institute,
Tanta branch

For Degree Of M.V.Sc In Veterinary Medical Science
(Bacteriology, Immunology and Mycology)

(2021)

جامعة بنها
كلية الطب البيطري بمشتهر
قسم البكتريا و المناعة و الفطريات

Benha university

Faculty of Veterinary Medicine

Department of Bacteriology ,

Mycology and Immunology

APPROVAL SHEET

This is to approve that the dissertation presented by Mohammed Mahmoud Mohamed Youssef wehaba to benha university entitled (the occurrence of a food pathogen in some types of soft cheese in the local market) for the degree of master (Bacteriology, Immunology and Mycology) has been approved by the examining committee 27/9/2020.

Committee members : **signature**

Prof.Dr/Ahmed Affifi Abd El ghafar Maarouf **(Chairman)**

Chief researchers institute manager

of animal health research institute

benha branch

Prof.Dr/Ashraf Awad Abd El tawab **(Member and supervisor)**

Professor of Bacteriology,Immunology

and Mycology Fac.of Vet.Med.

Benha university.

Prof.Dr/Amal Mostafa Eid **(Member and supervisor)**

Chief researchers of animal health

research institute benha branch.

Asso.prof.Dr/Fatma Ibrahim El-hofy **(Member)**

Assoc.Professor of Bacteriology,Immunology

and Mycology Fac.of Vet.Med.

Benha University.

Date /27/9/2020

سورة البقرة الآية: ٣٢

Acknowledgment

First of all, I wish to express my deep thanks to ***Allah***

Almighty who has endowed his help and blessings on me to have the present work achieved in such condition.

I wish to express my gratitude and thanks to ***Prof. Ashraf A wad Abdel El tawab*****,** *professor of Bacteriology, Immunology, Mycology, Faculty of Veterinary Medicine, Bennah University for his guidance, criticism, continuous encouragement, and helpful advice during this work.*

My thanks to ***Prof. Amal Mostafa Eid****, chief researcher23%,* ***Prof. Dalia Fathy Khater*** *, chief researcher87% , Animal Health Research Institute Tanta branch for thier encouragement and guidance during that work.*

Finally, I wish to express my warmest gratitude to all members of Bacteriology, Immunology, Mycology Dept., Faculty of Veterinary Medicine (bennah University) and Animal Health Research Institute members and all medical honest specialists.

Thanks, all for your patience and support.

Dedications

My Father

My Mother

My Sister

My Brother

My Uncles

My Family

Always all support religious forever

VITA

- *The Author Was Born On September 9, 1981, In El – Behera Governorate, Egypt.*

- *His Primary and Preparatory Educations Was Completed In 1993 In Saudi Arabia.*

- *His Secondary Educations Were Completed At Dalgmon Secondary School Village, Gharbia Province- Egypt - From Which Graduated In 1999.*

- *His Undergraduate And Professional Educations was Completed In Faculty Of Veterinary Medicine, Kefir El – Sheih University From Which He Received a BVSc. Degree In 2003. Very Good. The Seventh.*

- *He works at the Animal Health Research Institute, Tanta branch.*

LIST OF CONTENTS

Title	Page
1- Introduction	1-2
2- Review of Literature	3-19
1.2incidence of E.coli in soft cheese	3
2.2 phenotypic character of E.coli	13
3.2 in vitro antimicrobial sensitivity	13
4.2genotypic detection of virulence genes of E.coli	14
5.2 public health importance	19
3-Materials and methods	20-28
4- Results	29-34
5-Discussion	35-43
6- Conclusion and – Recommendations	44-45
7-Summary	46-47
8-References	48-68
9-Arabic summary	1

List of tables

Table No	title	Page No
1	Incidence of E.coli isolated from soft cheeses in El- Gharbia Governorate .	29
2	Regional Incidence of E.coli isolated from soft cheeses in El- Gharbia Governorate .	30
3	Biochemical tests	31
4	Results of serological identification of *E. coli* isolates.	31
5	Antimicrobials sensitivity results for *E. coli* isolates.	32
6	Results of molecular identification of *stx1* .*eae* A and *fim* H genes of *E. coli*	32

List of figures

Figure no	Title	Page no
1	Figure (1): Agarose gel electrophoresis of PCR amplified products of the virulence gene. For *Stx1* gene Were Negatives.	**33**
2	Figure (2): Agarose gel electrophoresis of PCR amplified products of the virulence gene. Lane L: DNA molecular size marker (100bP), lane Pos: Positive control, lane Neg: Negative control, Lane 1,2,3,4,5,6,7: for detections of *Fim H* virulence gene of *E.* coli. The size in base pairs (508bP) of PCR products is indicated for the bands. All seven isolates were positives.	**33**
3	Figure (3): Agarose gel electrophoresis of PCR amplified products of the virulence gene. Lane L: DNA molecular size marker (100bP), lane Pos: Positive control, lane Neg: Negative control, Lane 2,3: *Eae A* positives virulence genes of *E.* coli. The size in base pairs (248bP) of PCR products is indicated for the bands.	**34**

INTRODUCTION

1. INTRODUCTION

E. coli is one of the major significant bacteria, which has a bad effect on both human and animal species. Thus, this type of bacteria can deteriorate the milk particularly raw milk and other milk products as a result of poor hygienic measures (**Garbage *et al*., 2016; Lara *et al*., 2016).**

In Egypt Moreover, Kariesh cheese is a prevalent type of cheese that contains a higher protein content with a small number of fats (**Hamad, 2015).**

Damiati cheese is one of soft pickled cheese and makes up about 75% of the cheese produced and consumed due to its nutritional value, convenience and have a rich taste, strong sharp flavor as well as smooth body and texture.

Tallaga cheese is the most popular local type of packaged or unpackaged soft cheeses by all socioeconomic classes in Egypt due to its nutritional values, conveniences, and good tastes, *E. coli* is classified into six pathotypes: enteroaggregative, enterohemorrhagic Shiga toxin-producing *E. coli* (STEC), enteron invasive, enteropathogenic, enterotoxigenic, and diffuse adherent (**Jafari *et al*., 2012).**

Enteropathogenic *E. coli* (EPEC) is considered the most common bacteria isolated from cheese. EPEC also causes high morbidity and mortality rates among young and old people (**Kostas *et al*., 2010**).

In humans, the pathogenic effect of STEC nearly due to its ability to produce certain types of cytotoxins for example Shiga toxins (*stx1*), and intimin (*eae A*) virulent genes (**Slaney *et al*., 2009 and Assupol *et al*., 2015).**

Shiga toxins–producing *E. coli* (STEC), also known as neurotoxins-producing *E. coli* (VTEC) or Enterohaemorrhagic *E. coli* (EHEC), have been known as a group of highly pathogenic *E. coli* strains producing one or more Shiga toxins (**Monaghan *et al*., 2011).**

Escherichia coli (*E. coli*) is one of the normal inhabitant microorganisms of the large intestine in human and warm-blooded animals. The main source of *E. coli* in raw milk and milk products is fecal contamination together with poor hygienic practices **(Garbage *et al*., 2016; Lara *et al*., 2016).** However, certain serotypes of *E. coli* have acquired some virulence-associated genes that enable them to cause intestinal or Extra-intestinal diseases **(Hodson *et al.*, 2017).**

These serotypes that cause enteric infections are generally called diarrheagenic *E. coli* strains, and their pathogenesis is associated with a number of virulence attributes, which vary according to pathotype.

As soft cheeses (Kariesh, Tallaga, and Damietta) collected from different localities from the Gharbia governorate may be subjected to contaminations with *E. coli* and may constitute a public hazard. Therefore, this work was planned to covers the followings points:

1) Isolations and serologically identifications of pathologically pathotypes of Escherichia coli from Kariesh, Tillage, and Damietta types of cheeses in El _Gharbia governorate.

2) Molecular identifications of isolated bacteria (Bacterium Coli) using PCR Techniques and determination of some virulence genes from the isolated strains.

REVIEW OF LITERATURES

2- Review of literature

1.2 .incidence of E.coli in soft cheese

Alice *et al.* (2000) isolated *E. coli* colonies from 44 soft white cheese samples and EPEC serogroups O127 and O128 were found in 11.3% of cheese samples.

El-kosi (2001) examined bacteriologically the kareish cheese samples and *E. coli* recovered as 100 % of all samples.

Menendez *et al.* (2001) examined 24 of kareish cheese samples produced in Galicia. NW Spain for their microbiological characteristics. They found that *E. coli*.

Al – Jeddah and Robinson (2001) examined microbiologically white brined imported Damietta cheese samples from Egypt and purchased from food stores around Doha (Qatar) and found that all samples were free from *E. coli*.

Ahmed (2002*)* examined 20 samples of cheese for the presence *E. coli* and found that 25 % of the examined samples were contaminated with *E. coli*.

Gomez *et al.* (2002*)* examined bacteriologically 114 samples of soft cheese for the presence of *E. coli* and founded 0.9 % of samples were positive.

Amer and Ewin (2003) examined 40 random samples of street vendors kareish cheese collected from different localities in Alexandria and Kufr El-Sheikh governorates. *E. coli* could be isolated from 37.5% of the examined samples.

Thabit. S (2003) examined 40 samples of cheese for the presence of *E. coli* and found that 20% of the examined samples were contaminated with *E. coli*. the isolated strains were serotyped as O111, O119 and O157.

Spano *et al.* (2003) studied experimentally have already shown that treatment at high temperature such as curd stretching at 80 °C for 5 min during

the manufacture of cow's milk mozzarella cheese has been shown to effectively control STEC strain growth

El-Bakery (2004) examined bacteriologically 50 samples of Kareish cheese collected from different markets and shops from Giza, Egypt in summer and winter. He found that the percentage of *E. coli* in Kareish cheese was detected in 18 % and 26 in winter and summer respectively.

Mustafa (2004) found that 30.9% of the examined Damietta cheese samples were positive for *E. coli*.

Oxus *et al*. (2004) examined 50 white pickled cheese manufactured from raw milk for the presence of *E. coli O157 which* was detected in 4% of the total cheese samples.

Hunnish *et al.* (2005) reported that the number of cases deaths 13 in Canada with serotype O175: H7 of *E. coli* from ingestion of contaminated un pasteurized Gouda cheese.

Nemur (2005) isolated *E. coli* from Kareish cheese samples by ann incidence of 24% of the examined samples. The isolated serovars of *E. coli* were O111, O55, O126 and O157: H7 with a percentage 25%, 17 %, 33% and 25%, respectively.

Bah out and Mostafa (2006) examined 50 samples of Kareish cheese collected from local markets in Dakhlia province, Egypt *E. coli* were present.

Anand and Gangapur (2006) tested 77 samples of domestic soft cheese which were collected randomly from different groceries and markets in Kerman, Iran and they found that 98.70% of cheese samples were positive for *E. coli* and Eight *EPEC* serogroups were isolated from 15 (19.48%) of the samples.

EPEC serogroups included O26 (I), O86 (II), O114 (IV), O142 (IV and I), O119 (II), O127 (II), and O128 (III).

El-Pessary (2006) examined microbiologically Kareish cheese samples and pointed out those coliforms, fecal coliforms and *E. coli* were detected in 100%, 92.5% and 60% of the examined samples, respectively.

El-Pessary (2006) examined microbiologically Damietta cheese samples and pointed out *E. coli* were detected in 10% of the examined samples, respectively.

Mahi, Al-Ashmawy (2006) examined bacteriologically 25 samples of Kareish cheese collected from the El-Dahlia governorate for the presence of Coliforms and reported that coliforms were detected in all examined samples.

Johnson *et al.* (2006) found that *E. coli* causes illness ranging from gastrointestinal tract-related complications such as diarrhea, dysentery, urinary tract infection, pneumonia and even meningitis.

Anand and Gangapur (2006) tested 77 samples of domestic soft cheese which were collected randomly from different groceries and markets in Kerman, Iran and they found that 98.70% of cheese samples were positive for *E. coli* and Eight *EPEC* serogroups were isolated from 15 (19.48%) of the samples. EPEC servo groups included O26(I), O86(II), O114(IV), O142 (IV and I), O119 (II), O127(II), and O128(III).

Control and Prevention (206) weekly report which reported that during August-September 2005 a total of 52 (2.44%) of 2,160 inmates at a state correctional facility reported diarrhea, including 17 (33%) with bloody diarrhea. Sixteen of the samples were positive for the *Stx*1 gene but not for STEC O157 Serotyping, and in this case, the source of the Outbreaks likely was an ill food worker.

Laden and raze (2006) examined Seventy-seven random samples of domestic Iranian soft cheese that were collected from different groceries and markets in sterile conditions, cultured in selective media ., and examined for

biochemical tests. Among the 77 samples, *E .coli* was isolated from 76 (98.70%) of them and 15 (19.48%) of the isolates were EPEC. Eight EPEC identified serogroups were included: O26(I), O86(II), O114(IV), O142 (IV and I), O119(II), O128(III), and O127(II). All these types are pathogens. O127 was the most prevalent serogroup, followed by O128 and O119. The occurrence of a high proportion of *E. coli* in our cheese samples may be due to a lack of proper sanitation and the absence of pasteurization of milk used for cheese making. Therefore, strict hygienic measures must be followed and pasteurization of milk should be imposed to prevent contamination of cheese with coliforms, thus avoiding additional outbreaks of food-borne illness caused by *E. coli*.

Abraham *et al.* (2007) examined 80 samples of Kareish cheese collected from the Alexandria government. The prevalence of *E. coli* reached 75%, respectively.

Almeida *et al. (*2007) stated that out of 70 cheese samples were examined for the presence of *E.coli,* O157 for grading them as satisfactory, acceptable, unsatisfactory, or unacceptable and potentially hazardous; 22 were graded as satisfactory or acceptable, 37 were considered unsatisfactory because of the presence of *E.coli*, while 11 of the cheeses samples were graded as unacceptable and potentially hazardous because of the presence of excessive numbers of *E. coli,* in three of these samples.

Paneto *et al.* (2007) collected 50 samples of raw milk cheese from different supermarkets in Middle Western Brazil and were analyzed for *E. coli* which were recovered from 48 (96.0%) of the samples. The identified Serogroups were O125 (6.0%), O111 (4.0%), O55 (2.0%) and O119 (2.0%). Three (6.0%) and 1(2.0%) of *E. coli* isolates were verotoxigenic *E. coli* (VTEC) and enterotoxigenic *E. coli* (ETEC), respectively.

***Om*er *et al.* (2007)** analyzed 80 samples of kareish cheese collected from supermarkets shops and street vendors in Alexandria, Egypt for bacteriological

quality. The examined samples showed that *E. coli* was detected in 60 (75%) in Kareish cheese.

El-baradari e*t al.* (2007) examined microbiologically Damietta cheese samples and they found that 90% of the samples were positive for *E. coli*.

Nagaina and Chen (2008) Reported an association between virulence factors and antimicrobial resistance in *E. coli* strains isolated from dairy cows.

Osman and Omer (2008) found that *E. coli* was detected in some of the cheese samples.

Hassan (2008) reported that microbiological quality, composition, and some chemical properties were determined in 40 samples of Kareish cheese randomly collected from various markets located in Cairo. microbiological examination revealed that 12% of the examined kariesh cheese was contaminated with *E. coli.*

Sabry and Laila et *al*. (2008) examined bacteriologically a total of fifty raw milk and twenty-three Kareish cheese samples were randomly collected from different shops and distributors in Qena city *E .coli* were recovered from 11 (47.8%) of Kareish cheese samples.

Fadel *et al. (*2009*)* examined microbiologically 40 samples of Kareish cheese collected from supermarkets and dairy shops and street vendors in Ismailia city for the presence of *E. coli* and the results were 4 samples only were positive.

Fadel and Jaehn (2009) collected 40 Kareish cheese samples from different localities in Ismailia city and they could isolate 4 different *E. coli* serovars STEC group (O128: H_2), EPEC (O127), EIEC group (O124: H7) , and one untippable strain.

Ali *et al*. (2009) examined 15 random samples of Kareish cheese from dairy shops and street vendors in rural areas for Microbiological evaluation. The incidence of *E. coli* was 6.6% out of the examined samples.

Morass (2009*)* fifty-five samples of Kareish cheese were collected from different non inspected commercial establishments and submitted to the inspection of *E. coli* it was detected.

El-prince, Etnas *et al.* (2010) examined 30 random samples of Kareish cheese that were collected from a suit city, Egypt for fecal pollution indicators. All examined samples were contaminated with coliforms. The incidences of *E. coli* in Kareish cheese were 33.3%.

Kostas *et al.* *(*2010) cheese can be easily contaminated by food handlers who practice poor personal hygiene at some point during cheese preparation and/or assembly line.

Ortolani *et al*. (2010) studied the microbiological quality and safety of 18 soft cheese samples and they found that low levels of *E. coli* were observed.

Todd *et al.* (2010) several foodborne outbreak of colibacillosis investigation reports have identified the hands of food workers as the source of pathogens in the implicated food.

El Sayed *et al.* (2011) examined microbiologically (30 samples of Damietta cheese and 15 samples of Tillage cheese and 10 samples of Kareish cheese) were collected from Cairo and Giza governorates for the presence of *E. coli* and results respectively were (20%, 33,3%, 26.6%).

Ebrahim *Rahimi et al.* (2011) examined microbiologically 120 traditional fresh cheese samples for the presences of *Escherichia coli* in Iran state and the result was four isolates from traditional cheese and the genes *Stx1*

and *Stx2* were detected three *E. coli* isolates obtained from traditional cheeses samples.

Zeinhom (*2011)* tested 50 random samples of Kareish cheese were collected from different localities in Beni -Suef Governorate, Egypt monitoring of enteric pathogen with *E. coli* the data showed that *E. coli* was not isolated from any of the examined samples of Kareish cheese.

Hosni *et al.* (2011) made a survey to evaluate the level of microbial contamination of 35 samples of Kareish cheese retailed in the Cairo area and the results indicated that *E. coli* as a fecal indicator, was isolated from 25.7% of the total numbers of Kareish samples.

Ozan *et al.* (2011) studied the microbiological quality of white pickled cheese produced in small dairy plants which were investigated and found *E. coli* was isolated from 40% of the samples.

Dias *et al.* (2012) reported on the implementation of Good Manufacturing Practices **(GMP)** in a small processing unit of mozzarella cheese in Paraná, Brazil, and that the primary hurdles for the implementation of **GMP** were related to the difficulties to change the hygiene practices of food handlers

Ahmed (2012) examined microbiologically 25 samples of Damietta cheese and 25 samples of Talega cheese and 25 samples of Kareish cheese collected from dairy shops and street vendors in Beni Sufi city for the presence of *E. coli* and the result were 30%, 48%, 27.80%, and 39.94% respectively.

Abd El-Latifa, elan (2012) examined bacteriological 25 samples Kareish cheese randomly collected different farmers in El-Bahira, Egypt, and examined microbiologically. The author found that *E. coli* was detected in 44% in Kareish cheese.

Ahmed (2012) examined 25 random samples of Kareish cheese that were collected from different markets and retail shops in Beni -surf city, Egypt

for chemical and microbiological examination. The prevalence rate of *E. coli* isolation was 28.34%.

Arabi *et al.* (2012) investigated the frequency of *Fim H* and other adhesions genes in UPEC and determined the *Fim H* gene frequency as 87.7%.

Brooks *et al.* (2012) obtained 41 raw milk cheese from retail specialty shops, farmers' markets, and on-line sources, and the samples were then analyzed for the presence of *E. coli*. The results revealed that 2 of those contained *E. coli*.

Nagak *et al.* (2012) examined 120 random samples of raw milk, Damietta cheese, Kareish cheese, and Cooking butter (30samples each) which were collected from different localities in Assiut city, Egypt and they found that the incidence rates of *E. coli* in raw milk, Damietta cheese, and Kareish cheese samples were 83 %, 30%, and 60 %, respectively.

Samaha *et al*. (2012) examined 50 Kareish cheese samples that were collected from different localities in Alexandria and the results revealed that the rate of isolation of *E. coli* was 26%.

Clays *et al*. (2013) reported the occurrence of Pathogenic bacteria in raw milk and their products in pathogenic *E. coli,* Therefore, consumption of raw milk and traditional milk products can be caused foodborne pathogens Outbreaks.

Bunyanian *et al.* (2014) examined 50 samples of unpasteurized soft cheeses from retail shops which were analyzed for detection of *E. coli* and found 24 of the cheese samples were positive for *E. coli.*

Gundogan and Arci (2014*)* examined samples of white soft cheese from three different dairy-processing plants in Ankara City (Turkey) to find out if they were contaminated with *E. coli*. The results showed that the contamination percentages of *E. coli* were 60%.

Abo Zed (2014) examined microbiologically 30 samples of Kareish Cheese randomly collected from different dairy shops and supermarkets in Cairo, Giza, and Kilobit governorate, Egypt. The results revealed that *E. coli* failed to be detected.

El- kholy *et al.* (2014) examined bacteriologically 50 Fifty samples of Damietta Cheese were randomly collected from supermarkets, retailers, and shops in the Bani-Suif governorate, Egypt for the presence of *Escherichia coli*, and the result were 13 samples positive by percentage 26%.

Hussain *et al.* (2014*)* collected 120 samples of different dairy products including Cheese from different localities of Lahore and found 54% of tested samples were contaminated with *E. coli*.

Walid *et al.* (2014) examined bacteriologically 40 samples of Damietta Cheese and 20 samples of Kareish Cheese samples for the presence of *Escherichia coli* only 12 samples were positive for *E. coli* presence with 20 %.

Al Jabari *et al.* (2015) examined 70 samples of white Soft Cheese, the results revealed that the incidence of *E. coli* was 87%.

Aseel A. Saeed (2015) examined a total 50 Cheese samples were collected from a market in Al-Diwaniya City In Kuwait the study was done in the period between August and October 2014 and recorded 38 samples were positive for *E. co*li.

Met Wally and Fatma (2015) examined about 25 samples of Kareish Cheese collected from retailers and small dairy shops in Giza and Bani-Suief governorates markets for the presence of *E. coli* and result founded that about 72% of samples were positive for the *E. coli.*

Mohamed (2015) examined Kareish Cheese bacteriologically for the presence of *E. coli* and it is recorded from all samples by a percentage of 100 %.

Elnath *et al.* (2015) examined microbiologically 40 samples of Kareish Cheese samples were collected from different localities in El Monovia - governorate for the presence of *E. coli* and founded 50 % of samples were positive for pathogenic *E. coli.*

Gamal *et al.* (2015) examined microbiologically 50 samples of Damietta Cheese and 30 samples of Kareish cheese were collected from the Cairo governorate for the presence of *E. coli* and the results were 20% and 33% respectively.

El bigotry *et al.* (2015) examined microbiologically 40 samples of Kareish Cheese and 40 samples of Damietta cheese collected from supermarkets and dairy shops and street vendors in El- Gharbia governorate for the presence *E. coli* and the results were 52.5 % and 15% respectively.

Jehan e*t al.* (2015) examined microbiologically 30 samples of Kareish Cheese and 30 samples of Damietta cheese from different shops and supermarkets and street vendors in Ismailia city for the presence of *E. coli* and the results were 9 samples of Kareish cheese were positive and 12 samples of Damietta Cheese were positives.

Abo Baker *et al.* (2016) isolated from fresh locally made Soft Cheeses were 9 (33 %) and 3 (11 %) respectively from (masoor and Ricotta soft cheeses) positives for the presence of *E. coli.*

Gamal and soda (2016) examined 50 samples of Damietta cheese and 50 samples Kareish Cheese in Cairo and Giza governorates for the presence of the *E.coli.*

2.2 .phenotypic character of E.coli

Viipuri *et al.* (2013*)* examined 50 samples of soft cheese for the presence of *E. Coli* and founded that 28% of samples were positive for *E. coli* were collected from milk vendors, retail shops located in Anand city, (India) under aseptic precautions. For the enrichment of the organism from the collected samples, MacConkey broth was used and inoculation was carried out on MacConkey agar and EMB agar. Later on, to confirm the isolates, various biochemical tests such as the IMIS test, Urease test were performed. Evaluation of the antibiotic sensitivity pattern of *E. coli* was assessed by the disk diffusion method. Finally, the *E. coli* isolates were screened for the presence of virulence - associated genes by PCR and founded that 7 isolates of *E. coli* positive for the presence of *Stx1* (8.75 %) only and 5 isolates were positive for The presences of both *Stx1* and *Stx2* and 7 isolates were positive for *Eae A* gene.

Lotfy *et al*. (2017) examined 25 samples of each Kareish cheese and Damiatte cheese collected from Mansoura city for the presence of *E. coli* and founded the results were 32% and 16% by using MacConkey agar medium and 44% and 20 % by using EMB medium.

3.2 .in vitro antimicrobial sensitivity

Cergol-Novella *et al.* (2011) showed that the highest resistance rates of STEC were identified for tetracycline (100%), streptomycin (78%), and trimethoprim-sulfamethoxazole (56%). Furthermore, all strains were harbored *Stx* genes.

Assupol *et al.* (2015) demonstrated that the STEC isolates that carried *Stx1* only showed the highest resistance to Streptomycin

(Pure Negative) (85.6%), followed by Kanamycin (72.2%), and Nalidixic acid (70.0%).

4.2. genotypic detection of virulence genes of E. coli

Pradel *et al.* (2000) reported the isolation of STEC non-O157 strains from cheese in France, all of them carrying the *stx*1 gene but not the *eae A* gene.

Rey *et al*. (2006) showed that 3 STEC carried the *Stx1* gene, 5 possessed the *Stx2* gene, and 1 both *stx1* and *Stx2*. Whereas only O157: H7 serotype showed *eae A* virulence gene.

Lothar Boutin *et al.* (2007) examined 219 Shiga toxins producing *Escherichia coli* (STEC) strains from meat, milk, and cheeses samples collected in Germany between 2005 and 2006 and they found that Shiga toxin *Stx1* present in 88 (40 %) strains.

Garofalo *et al.* (2007) studied 18 UPEC isolates collected from females and found that the *Fim H* gene was the most prevalent virulence factor and 100% of the isolates had that gene.

Irion *et al. (*2007) it is also important to mention that in Brazil, human STEC infections have been primarily associated with diarrhea and most of the STEC strains isolated are non-O157 strains harboring *Stx*1 and *Eae A* genes.

Malden *et al. (*2009) evaluated the distribution of virulence genes among the studied UPEC and showed that the prevalence of different virulence genes varied from 10% for the CNF colony necrotizing factor gene to 80% for the *Fim H* gene**.**

Watts *et al.* (2010) demonstrated that the *Fim H* gene was the most frequent virulence gene and was detected in 98% of *E. coli* strains isolated from patients with UTIs.

Shaath Thanom Ahmed *et al.* (2011) examined microbiologically 115 samples of white cheeses from 3 different areas in Baghdad surveyed and examined for the presence of *Escherichia coli* the result showed 7 samples are

positive for the presence of *Escherichia coli* and one strain contain *Stx1* virulence genes.

Mohammed Elhadidy and Mahmoud hammed (2012) examined microbiologically 124 raw milk cheese samples (64 Kareish and 60 Damietta cheese samples) for the presences of STEC using molecular detection of virulence markers such as Shiga toxin *(Stx1)* and intimin gene *(Eae A) not far enough* and by Serotyping they found only 14 samples positive for the *E .coli*.

Elhadidy and Mohammed (2013) studied microbiologically 124 raw milk cheese samples (Kareish and Damietta cheese samples) for *E. coli* contamination. They found that 14 *E. coli* strains were positive for STEC. **Elhadidy and Mohammed (2013)** reported that STEC incidence in 124 cheese samples was 11.29%. All recovered isolates harbored the *Stx2* gene 100% either single or in association with the *Stx1* gene, while *Stx1* was present in 64.28% of isolates. They reported that theintimin gene (*Eae A*) was present in 21% of isolates.

Tarkhun *et al.* (2013) reported that among the studied virulence genes of UPEC strains, the *Fim H* gene was the most prevalent virulence gene and was founded in 68% (61/90) of the UTI isolates.

Durra & Marin (2014) examined 330 *E. coli* strains isolated from 111 Minas Fresca cheeses from Southwest Minas Gerais State and also did not detect *Stx* 1, *Stx* 2, or *Eae A* genes among the *E. coli* isolates.

Hoffmann *et al.* (2014) analyzed 87 *E. coli* strains isolated from pasteurized milk obtained from 22 farm dairies located at Parana state in Brazil and did not detect the *Stx*1, *Stx*2, and *Eae A* virulence genes among the *E. coli* isolates.

Assupol *et al. (*2015) reported from 561 *E. coli* isolates, 90 (16.0%) were carriers of *Stx1*, 97 (17.3%) of *Stx2*, and 45 (8.0%) of *Eae A* genes singly. 37

(6.6%) isolates were carriers of *Stx1* and *Stx2*, 110 (19.6%) were carriers of *stx1* and *Eae A*, and 67 (11.9%) were carriers of *Stx2* and *Eae A.*

Zahren Hejazi *et al.* (2015) a total of 140 isolated *E. coli* strains from patients with UTI were identified using biochemical tests and then evaluated for the *Fim H* gene by polymerase chain reaction (PCR) analysis. The *Fim H* gene was found in 130 isolates (92.8%) of the UPEC strains (How Many Isolates and strains There are big Differences). 130 isolates positives for the *Fim H* gene, 62 (47.7%), and 68 (52.3%) belonged to hospitalized patients and outpatients, respectively.

The results of this study indicated that more than 90% of *E. coli* isolates harbored the *Fim H* gene . The high binding ability of *Fim H* could result in the increased pathogenicity of *E. coli*; thus, *Fim H* could be used as a possible diagnostic marker and/or vaccine candidate.

Mubarak *et al.* (2016) examined 55 Kareish Cheese samples for the presence of *E. coli* and found that 41 samples (74.4%) were positives for *E. coli* Occurrences and these isolates screened by PCR for the presences of virulence genes *Stx1* and *Eae A* virulence genes and the percentages were (90 %), (45%) respectively.

coli and they founded that the results were negatives.

Nabil *et al.* (2016) examined 95 food and environmental samples were examined for the presence of Shiga Toxigenic *Escherichia coli* for the presences of *Stx1* and founded one sample positive for the presence of STEC.

Same a wad (2016) examined microbiologically 15 samples of traditional Kareish Cheese made using raw milk were collected from different markets in different governorates (Alexandria, Behera, Kefir el- Shiekh, Damietta, Gharbia) for the presences of *E. coli* and the results were negatives.

Furthermore, the results showed that 11 out of 11 (100%) *E. coli* had both *Stx1* and *Stx2* while *Eae A* gene was not found in *E. coli* isolated from traditional milk products. The *Eae A* gene is an accessory virulence factor for STEC that is thought to enhance the virulence of STEC, while some STEC strains not harboring *Eae A* gene have been shown to cause human illnesses.

Cholesteric *et al.* (2017) examined microbiologically 26 Kind Cheeses samples randomly purchased from local markets in Kashan, Iran and evaluated for the presence and Occurrence of STEC by culturing and Molecular methods and the incidence of *E. coli* in 26 samples was as follow 11,54 % furthermore isolates were assayed for the presences of Shiga toxins *Stx1* and intimin gene *eae A* . the findings showed that 100 % of *E. coli* contain *Stx1* while *Eae A* gene was no found in *E. coli* isolates of traditional milk products In recent years, Shiga toxigenic *E. coli* (STEC) strains as emerging groups of foodborne pathogens are responsible for most foodborne illnesses. The pathogenicity of these strains is almost related to the ability of the microorganism to produce the one or two cytotoxins called Shiga toxins (*Stx1* and *Stx2*), and the presence of the intimin (*Eae A*) gene, which is responsible for intimate attachment of the organism to the intestinal epithelium.

Douellou *et al.* (2017) demonstrated that the virulence genes profiles of dairy products and human STEC strains were similar to **Farhad shaft chaleshtori *et al.* (2017)**.

Patricia Cardoza and Jose miacid marine (2017) Examined 59 raw milk Cheese samples collected from local producers in Brazil for microbiological analysis and found that 38 (64,4%) tested positive for *E. coli.* then 147 strains of

E. coli from positives samples and screened by polymerases chains reactions for the presence of virulence genes *Stx1* (verotoxin type 1) and *Eae A* (intimin) gene and found that (9,5 %) were positives for the presence the *Stx1* gene and (11) of

them also harbored *Eae A* gene. all STEC – contaminated food can be considered potentially hazardous.

Hussein *et al.* (2019), inspected 75 samples of Kareish Cheese collected from different areas presented in Monofia- governorate According to the results, the frequency of *E. coli* isolated from Kareish was 16%. Serological identification classified the *E. coli* strains into three groups: Enterohemorrhagic *E. coli* (EHEC) serogroups (O26: H11, O91: H21, O111: H2 ,and O103: H2) While the Enterotoxigenic *E. coli* (ETEC) Serogroups were detected as O125: H21 which is the most prevalent strain. O171: H2, O86, and O119: H6 belonging to Enteropathogenic *E. coli* (EPEC). The most prevalent genes detected in *E. coli* strains were *Stx1* (87.5), *Fim H* (75), and *Eae A* (25%) genes. In conclusion, Kareish Cheese is extremely tainted with pathogenic *E. coli* strains, which represents a strong hazard on the Animals and Human health.

Taha *et al.* (2019) examined microbiologically 90 Cheese samples high salt soft cheese, Kareish Cheese, and Talega cheeses were examined for *E. coli*. Serodiagnosis of *E. coli* has been done by slide agglutination test. Toxigenic *E. coli* isolates were detected using PCR assay. Results postulated that the detection rate of non-O 157 was 36,33% in Kareish Cheeses, 20 % in high salt soft Cheeses, and 33,33% in Talega cheeses. Ten different serotypes of *E. coli* non O157 Serotypes *of E. coli* non-O 157 have been identified as follows: O1, O18, O20, O25, O26, O125, O126, O127, and untyped *E. coli*. regarding PCR results: all serogroup was negative for the *Stx1* gene, only servo group 25 positives for *Eae A* gene.

Ibrahim *et al*. (2019) examined bacteriologically 30 samples of Damietta cheese and 30 samples of Kareish Cheeses randomly collected from local markets, farmers vendors, and supermarkets at different localities in the Nile Delta region, Minufia- governorate, Egypt. *E. coli* recovered from samples were 13 (43,3 %) and 22 (73.3 %) respectively. The Serotyping reveals in Kareish

Cheeses as following O128; H2, O26; H11, O44: H 18, O119: H6, O127: H6, O127: H6, O111: H2, O124, O146: H21 and in Damietta cheese as following :O128: H2, O26: H11, O91: H21, O127: H6 , O111: H2, O124, O146 H41and PCR results as following: O26; H11 had *Stx1* and *Eae A* genes, O44: H 18 had not S*tx1* and *Eae A* gens, O91: H21 had *stx1* and had not *Eae A*, O111: H2 had *stx1* and *Eae A* genes, O124 had not *stx1* and *Eae A* genes.

5.2. public health importance

Hussein and Sakuma (2005) studied the role of dairy cattle and their products in human infections with Shiga toxin-producing *Escherichia coli* (STEC). Shiga toxin-producing *E. coli* were isolated from raw milk, milk filters, and dairy products such as yogurt and soft cheeses. STEC infection was attributed to the consumption of dairy products that were contaminated with cattle feces. Thus, dairy cattle are considered reservoirs of STEC and can impose a significant health risk to humans.

Espies *et al.* (2006) reported that there were many cases of deaths in France due to the contamination of fresh unpasteurized goats' cheese by *E. coli* serotype O157: H7**.**

Ibrahim *et al*. *et al.* (2020) examined non-O 157:H 7 Shiga-toxin producing *Escherichia coli* (STEC) is a major foodborne pathogen that causes severe disease in humans ranged from mild gastroenteritis to critical illness and death especially O 26. O 111 in kareish cheeses**.**

MATERIALS AND METHODS

3- Materials and Methods

1. Materials

3.1. Samples:

One Hundred and fifty random samples of soft cheeses (50 samples of Kariesh Cheese – 50 Samples Of Tallaga Cheese – 50 Samples Of Domiatti Cheese) Were Collected Randomly From Small Diaries, Local Super Markets, Farmers Houses, Shops ,And Street Vendors In Different Localities At Gharbia Governorate. The Collected samples were transferred directly to the laboratory in an ice – box under aseptic conditions without delay and then subjected to bacteriological examination.

3.2.1. Isolations of *E*. coli:

5 grams of samples of each (Kariesh Samples – Tallaga Cheese- Domiatti Cheese) were added to 25 ml sod. Citrate and blended for two minutes for the examination of *E. coli* by these tests. (Preparation of samples): (maintain PH) (flushing)

3-2-1.1- MacConkey's broth test

(Acid And Gas Test): Two ml of each blended samples are added to MacConkey broth with bin Dorf tubes and incubated for24 hours at 37^{o}C for the detection of acid and gas if it is presently indicated for the presence of *E. coli.* (Selective Media is Gram -Negative Facultative Anaerobic Enterobacteriaceae)

3-2-1.2- Eosin Methylene Blue Media Test:

From the positive MacConkey's broth test with yellow colorations and acid and gas results are cultured on the Eosin Methylene Blue agar plates (EMB) is Selective Media for *E. **coli,*** Then Incubated at 37 ºC for 24 hours. Typical colonies of *E. coli* appeared Greenish Metallic with or Without Dark Black Centers. Suspected colonies were picked up and transferred to nutrient agar

slopes and then incubated at 37 ºC for 24 hours. The purified colonies were subjected to Biochemical and Serological Examinations.

3-2-2: - Biochemicals Identifications:

The pure colonies of isolates were identified biochemically as fellow: -

3.2.2.1- Indole Test: (Positive)

To each 48hrs incubated culture at 37ºC in 1% peptone water, one ml of ether was added. The tubes were vigorously mixed and allowed to stand for a few minutes until the ether rises to the surface. About 0.5ml of Kovac's reagent was added on the side of the tube gently. The Development of a rosy color ring indicated the presence of indole.

3.2.2.2- Citrate Utilization Test: (Negative)

The ability to utilize citrate was detected by making a single streak over the surface of Simmons's citrate agar slope, then incubated at 37ºC for 7days. The development of blue coloration indicated citrate utilizations.

3.2.3-Serological Identifications of *E. coli* serovars:

Two separate drops of saline were put on a glass slide and a portion of the colony from the suspected culture was emulsified with the saline solution to give a smooth fairly dense suspension.

- To one suspension, control, one loopful of saline was added and mixed. To the other suspension, one loopful of the undiluted antiserum was added and titled back and forward for one minute.
- Agglutination was observed using indirect lighting over a dark background. When a colony gave a strongly positive agglutination with one of the pools of polyvalent serum, a further portion of it was inoculated onto a nutrient agar slant and incubated at 37C for 24 hours to grow as a culture for testing with monovalent sera.

- A heavy suspension of bacteria from each slope culture was prepared in saline, and slide agglutination tests were performed with the diagnostic sera to identify the O-antigen.

3.2.4-In-Vitro antibiotic sensitivity of E. coli:

Subcultures from the isolates were prepared and the test was applied as follows:

A smooth single colony was inoculated in 5 ml nutrient broth and incubated at 37°C for 18 hrs., then turbidity was adjusted to 0.5 McFarland contain (1.5×10^8) colony-forming unit/ml, then few drops of the inoculated broth were flooded on to the surface of Muller-Hinton agar plates. Excess of cultural fluid was removed aseptically and the plates were allowed to stand for 15 minutes at 37°C for dryness. Then the inoculated plates were overlaid with antibiotic discs using sterile forceps, the discs were distributed in a manner where the distance among them was optimum and away from the edge of the plate to avoid overlapping of inhibition zones and give more wide area for the zone of inhibition. The inoculated plates were incubated at 37°C for 24 hours. The Inhibition zones, in mm, were measured and scored as sensitive, intermediate, and resistant categories with the critical breakpoints recommend via CLSI **(CLSI, 2018).**

3.2.5-Polymerase Chains Reactions Techniques (PCR):

3.2.5.1- Materials used for extractions of DNAs:

3.2.5.1.1-QIAamp DNA Mini Kit-Catalogue no.51304.

The QI amp DNA Mini Kit provides silica-membrane-based nucleic acid purification from different types of samples. The spin-column procedure did not require mechanical homogenization, so the total hands-on preparations time is only 20 minutes.

3.2.5.1.1-Ethanol 96% Apply chem.

3.2.5.2-Equipments and apparatuses used for extractions of nucleic acids:

3.2.5.2.1-Epindorff tubes 1.5 ml capacity.

3.2.5.2.2--Monochannel micropipettes 20-200 µl, 100-1000 µl (Biohit).

3.2.5.2.3--Sterile filter tips. (200 µl,1000 µl) capacity.

3.2.5.2.4-Centrifuge (Sigma sartorius).

3.2.5.2.5-Type II-A biosafety cabinet (Thermo).

2.5.3-PCR Master Mix used for c PCR: -

3.2.5.3.1-Emerald Amp GT PCR master mix (Takara)Code No. RR310AContains:

a) Emerald Amps separates the GT PCR master mix (2x premix).

b) PCR grade water

c) Oligonucleotides primers used in PCR.

They have specific sequences and amplify a specific product as shown in Table (primers-securities imported bio keys).

Table (specific not copied): Oligonucleotide primers sequences

Source: Metabolon (Germany).

Selected V Genes	Sequences	Amplified products	References
St 1	ACACTGGATGATCTCAGTGG	614 bp	Dipineto et al., 2006
	CTGAATCCCCCTCCATTATG		
Fim H	TGCAGAACGGATAAGCCGTGG	508 bp	Ghanbarpour and Salehi, 2010
	GCAGTCACCTGCCCTCCGGTA		
Eae A	ATGCTTAGTGCTGGTTTAGG	248 bp	Bisi-Johnson et al., 2011
	GCCTTCATCATTTCGCTTTC		

3.2.5.4-DNA Molecular weights markers: -

3.2.5.4.1-Gel Pilot 100 bp plus ladder (cat. no. 239045): - supplied from QIA GEN (USA)

A) - Number of bands: 11

B) - Sizes ranges : 100-1500 bp.

C) - Genes rulers : 100 bp DNA ladder (cat. no. SM0243) supplied from ferments.

3.2.5.5-Materials used for agarose gels electrophoresis:

3.2.5.5.1- Agarose 1.5% (Sambrook *et al.*, 1989)

A multi-purpose, high gels strengths agarose is suitable for a wide range of molecular biologics techniques. It has high gel strength and limited differentiated limits, multi Agarose could effectively separate large DNA fragments with reduced running times. This in turn means less band diffusion, a problem often associated with long running times.

It was prepared as follow:

A) - Agarose powders (AP genes): 1.5 g.

B) - TBE : 100 ml.

D) - Ethidium bromide solution 10 mg / ml (Sambrook *et al*., 1989)

i) - Ethidium bromide powders (Sigma) 10 mg.

ii) - Sterile DW 1.0 ml.

It was mixed and stored covered at 4˚C

It was added to melted agarose to reach a final concentration of 0.1-0.5 μg/ml.

C) - Tris borate EDTA (TBE) electrophoresis buffer (1x) (WHO, 2002) :

i) - Tris buffer (Fluka) : 10.78 g.

ii) - Boric acid (Fluka) : 5.5 g.

iii) - EDTA for DNA Fragments easily separations (Winlab) : 0.82 g .

It was brought up to 1 liter with deionized water, pH was checked up. If the pH was out of the range of (8-8.6), new solutions were surely preparing.

Any changes in ions concentrations will affect the migrations of the DNA Fragments through the gel.

3.2.5.6-Equipment and apparatuses used in PCR :-

i)- Calibrated cylinders.

ii)- Glass flasks.

iii)- CR tubes 0.2 ml capacity.

iiii)- (Scales Tec).

iv)- microwave (Panasonic).

v)- mono channels micropipettes (2-20 µl) (Bio hit).

vi)- Sterile filter tips.

vii)- Gel casting apparatus (Biometry).

viii)- T3 Thermal cycler (Biometry).

viiii)- power supply (Biometry).

viiii)- Type II A biosafety cabinet. (Thermo).

x)- Gel documentation system. (Alpha Innu tech).

xi)- Deionizer (Millipore).

xii)- Double distillatory (Sanyo).

3.2.5.7-Method:-

A- Extractions of DNAs according to QI amp DNA mini kits instructions:

1. rpm for 1 min. The QI amp mini spin column was placed in a clean 2 ml collection tube, and the tube containing the filtrate was discarded.
2. The QI amp mini spin column was carefully opened and 500 ml buffer AW1 was added without wetting the rim. The cap was closed, and centrifugated at 8000 rpm for 1 min. The QI amp mini spin column was placed in a clean 2 ml collection tube, and the tube containing the filtrate was discarded.
3. The QI amp mini spin column was carefully opened and 500 ml buffer AW2 we added without wetting the rim. The cap was closed, and centrifugated at full speed for 3 min.
4. The QI amp mini spin column was placed in a new 2 ml collection tube and the old collection tube was discarded with the filtrate. Centrifugation at full speed for 1 min was done.
5. The QI amp mini spin column was placed in a clean 1.5 ml microcentrifuge tube, and the collection tube containing the filtrate was discarded. The QI amp mini spin column was carefully opened and 100 µl buffer AE was added. The QI amp mini spin column was Incubated at room temperature (15-25˚C) for 1 min, and then centrifugated at 8000 rpm for 1 min.

2- Preparations of PCR (s) Master Mix according to Emerald Amp GT PCR master mix (Takara): Code No. RR 310A kit.

Component	Volume/reaction
Emerald Amp GT PCR master mix (2x premix)	12.5µl
PCR grade water	4.5 µl
Forward primer (20 pmol)	1 µl each
Reverse primer (20 pmol)	1 µl each
Template DNA	6 µl
Total	25 µl

3- Cycling conditions of the primers during PCR.

1. The Temperature and time conditions of the two primers during PCR are shown in Table (paid specifics) according to specific authors and Emerald Amp GT PCR master mix (Takara) kit.

Showing Tables: Cycling conditions of the different primers during

PCR Techniques: -

Genes	Primary denaturation	Secondary denaturation	Annealing	Extension	No. of cycles	Final extension
Stx1	94˚C 5 min.	94˚C 30 sec.	58˚C 40 sec.	72˚C 45 sec.	35	72˚C 10 min.
FimH	94˚C 5 min.	94˚C 30 sec.	50˚C 40 sec.	72˚C 45 sec.	35	72˚C 10 min.
EaeA	94˚C 5 min.	94˚C 30 sec.	51˚C 30 sec.	72˚C 30 sec.	35	72˚C 7 min.

4- DNAs Molecular weights markers: -

The ladder was mixed gently by pipetting up and down. 6 µl of the required ladder were directly loaded.

3-5- Agarose gel electrophoreses (Sambrook *et al*., 1989):-

Electrophoresis grade agarose (1.5 g) was prepared in 100 ml TBE buffer in a sterile flask, it was heated in a microwave to dissolve all granules with agitation, and allowed to cool at 70˚C, then 0.5µg/ml ethidium bromide was added and mixed thoroughly.

The warm agarose was poured directly in the gel casting apparatus with the desired comb in the apposition and left at room temperature for polymerization.

The comb was then removed, and the electrophoresis tank was filled with TBE buffer. Twenty µl of each unity plex PCR product and 30 µl of each multiplex PCR product, negative control, and positive control were loaded to the gel. The power supply was 1-5 volts/cm of the tank length. The run was stopped after about 30 min and the gel was transferred to the UV cabinet. The gel was photographed by a gel documentation system and the data we are analyzing's through computers soft wares.

Results

4- RESULTS

The morphological characters of the colonies on Eosin Methylene Blue were a brilliant green. A total of 64 bacterial isolates were obtained from examined samples, 27 Bacterial Isolates From Kareish Cheeses and 21 Bacterial Isolates From Damietta Cheeses And 16 bacterial isolates from Tallaga Cheese *E*. coli Isolated From Kareish Cheeses With An Incidences (54%) *E*. coli isolated from Damietta cheeses with an incidence (42%) *E*. coli isolated from Tallaga cheese with an incidence (32%) as shown in (Table 3).

E. coli gives Positive results to Indole Test results and Simmons citrate Negative.

Sex *E*. coli isolates were serotyped, Five from Kareish Cheeses, and One from Tallaga Cheese, Serological Identification revealed that Five isolates from Kareish Cheeses were belonging to (O119: H7, O111: H8, O86: O18: H1: K7 and O128). One isolates from Tallaga Cheese belonging to (O124).

Table (1): Incidence of *E. coli* isolated from soft cheeses in El-Gharbia - Egypt:

Type Of Sample	Number of The Sample	Number of Positive samples	Percentage of Positive samples
Soft Cheese	150	64	42.5 %
Kariesh Cheese	50	27	54%
Tallage Cheese	50	16	32%
Domiatti Cheese	50	21	42%

Tables 2. Regional Incidence of E.coli in Soft Cheese :

Regions/Kariesh Cheese	Number	Positive Results	Percentage
Kafer EL-Zayate City	10	2	20 %
EL-Santta	10	Zero	Zero
EL-Mahlla EL-Cobra	10	10	100 %
Kottor	10	10	100%
Tanta	10	5	50 %

Regions/DomiattCheese	Number	Positive Results	Percentage
Kafer EL-Zayate City	10	Zero	Zero
EL-Santta	10	10	100 %
EL-Mahalla EL-Cobra	10	5	50%
Kottor	10	5	50 %
Tanta	10	1	10

Regions/ TallagaCheese	Number	Positive Results	Percentage
Kafer EL-Zayate City	10	ZERO	ZERO
EL-Santta	10	ZERO	ZERO
EL-Mahlla EL-Cora	10	4	40%
Kottor	10	7	70 %
Tanta	10	5	50 %

Table (3) Biochemical tests of *E*. coli:

Bacteria	Biochemical test	Biochemical test
	Indole	Citrate utilization
E. coli	+	-

Table (4): Results of serological identification of *E*. coli isolates: -

Serial	Identified Bacterium	Polyvalent	Monovalent	Strain
1	*E.* coli (K3)	1	O119	EPEC
2	*E.* coli (K1)	1	O111	EHEC
3	*E.* coli (K2)	1	O86	EPEC
4	*E.* coli (K5)	3	O18	Ex PEC
5	*E.* coli (K6)	1	O128	EIEC
6	*E.* coli (D3)	7	O124	EPEC

K =Kariesh Cheese, D=Domiatti Cheese , T= Tallaga Cheese.

Table (5): Antimicrobials sensitivity results for *E.* coli isolates according to CLSI (2018):

Antimicrobial agent	Diffusion zone break point (mm)	E.coli D3 O124	E.coli K3O119	E.coli K2 O186	E.coli K1 O111	E.coli K5O18	Sensitivity Percent
Ciprofloxacin (CIP)	12 (mm)	S(20)	S(28)	S(29)	S(27)	S(32)	100%
Enrofloxacin (ENR)	0.5(mm)	S(8)	S(9)	R(0.3)	S(7)	S(8)	83.33%
Chloramphenicol (C)	15(mm)	S(23)	S(26)	R(7)	R(8)	I(17)	59%
Oxytetracycline (OT)	14(mm)	S(22)	S(21)	R(11)	S(25)	S(24)	83.33%
Neomycin (N)	12(mm)	I(14)	R(9)	R(8)	R(9)	R(7)	%8
Nitrofurantoin (F)	32(mm)	R(25)	S(75)	R(24)	R(19)	S(95)	50%
Erythromycin (E)	13(mm)	R(9)	I(17)	R(9)	S(28)	R(7)	25%

S= sensitive R= Resistant I = intermediate

K= Kariesh cheese D=Domiatti cheese

Table (6): Results of molecular identification of *stx1, eae A* ,and *fim H* genes of *E.* coli:

Sample	*eaeA*	*fimH*	*stx1*
1	-	+	-
2	-	+	-
3	+	+	-
4	+	+	-
5	-	+	-
6	-	+	-
7	-	+	-



Figure (1)

Agarose gel electrophoresis of PCR amplified products of the virulence gene. For *Stx1* genes were Negatives.



Figure (2)

Agarose gel electrophoresis of PCR amplified products of the virulence gene. Lane L: DNA molecular size marker (100bP), lane Pos: Positive control, lane Neg: Negative control, Lane 1,2,3,4,5,6,7: for detections of *Fim H* virulence gene of *E.* coli. The size in base pairs (508bP) of PCR products is indicated for the bands. All seven isolates were positives.



Figure (3)

Agarose gel electrophoresis of PCR amplified products of the virulence gene. Lane L: DNA molecular size marker (100bP), lane Pos: Positive control, lane Neg: Negative control, Lane 2,3: *Eae A* positives virulence genes of *E.* coli. The size in base pairs (248bP) of PCR products is indicated for the bands.

DISCUSSIONS

5- Discussions

E. coli has largely been regarded as an opportunistic pathogen causing both intestinal and extraintestinal diseases associated with large numbers of food poisoning outbreaks **(Varnam and Evans, 1991*).***

Enteropathogenic *E. coli* causes about 30-40 % of infantile diarrhea, as it is the most frequently isolated bacterial pathogen from the cheeses ***(*Doyle and padyhe, 1989)**.

Escherichia coli is commonly present in the intestinal tract of humans and animals, and its detection in processed foods is an indication of fecal matter contamination. It also indicates the inadequate control of the environment of the cheese processing unit, the personal hygiene of the food handlers, and the poor quality of the finished product **(Kostas *et al.*, 2010*;* Dias et *al.*, 2012*).*** Because the occurrence of an outbreak of foodborne disease is caused by enteropathogenic *E. coli* and is associated with the consumption of soft - ripened cheese ***(*Maurier *et al.*, 1973),*** the presence of these microorganisms in cheese has taken on added significance.

E. coli* is** responsible for several outbreaks of diarrhea in children and adults after ingestion of contaminated milk and dairy products. Different studies showed that 1 - 5% of food-borne infections were related to consumption of milk and dairy products, that 53% of cases of food-borne infections caused by contaminated cheese , and that enteropathogenic ***E. coli (EPEC) is the causative agent of 18.33% of these cases ***(*Schade *and* Yeager, *2001).***

EPEC strains have been implicated in food-borne human illnesses, especially as an important agent of infantile diarrhea in developing countries ***(*Silva *et al.*, 2001)**. EPEC strains are genetically mutative and cause different types of diarrhea in patients. A combination of these strains with

enterohemorrhagic *E. coli* increases their ability to create systematic infections in humans (**Desilk *et al.*, 1991).** The pathogenicity of EPEC strains is not known completely but most of them produce Vero toxins which are different from the enterotoxigenic *E. coli* (ETEC) toxins ***(*Campos *et al*., 1994*;* Crossly *et al.*, 1955).** Some EPEC strains are attached to the intestine epithelium ***(Campos et al.*, 1994).** Different strains of EPEC cause a different level of infection, and it has been shown that these strains are continuously changing and their genome becomes similar to hemorrhagic strains ***(*Campos *et al.*, 1994).**

Coliforms could be found in cheeses and are used as a hygienic indicator for such products. The presence of coliforms in cheese and their relation to enteropathogenic ***E. coli*** in soft cheeses have received considerable attention in previous studies ***(*scared *and* Yeager, 2001; Maurier *et al.*, 1973*).*** The relation between pathogenicity and different serotypes of ***E. coli*** has been suggested and proved ***(*Crossly *et al*., 1995*).***

Contamination of dairy products by EPEC strains has been investigated and O126, O128, O25, and O125 were isolated (**Abbar, 1988; Ahmed *et al.*, 1988*).*** Cheese has been known as an important animal origin product and in most countries, domestic and traditional produced cheese is in demand.

Shiga toxin-producing ***E. coli*** (STEC) are among the most important causes of foodborne diseases. They are responsible for several human gastrointestinal diseases, including watery or bloody diarrhea (**Paton & Paton, 1998).** The primary virulence factor of STEC is the production of common feature and the Shiga toxin 1 (Stx1), Shiga toxin 2 (Stx2) , or its variants, or a combination of these. Highly human-virulent STEC strains are also called enterohemorrhagic ***E. coli*** strains (**Bolton, 2011).**

Food is an important vehicle for the transmission of STEC strains, and most of the foodborne outbreaks caused by STEC have been associated with

unpasteurized milk and its derivatives **(Farrokh *et al.*, 2013).** This study aimed to assess the STEC occurrence in the collected cheeses samples.

Pathogenic STEC strains usually contain other virulence factors such as the outer membrane protein intimin (*Eae* gene), a protein essential for the intimate attachment, and the formation of attaching and effacing lesions on gastrointestinal epithelial cells (**Bolton, 2011*).*** Most outbreaks and sporadic cases of hemorrhagic colitis (HC) and hemolytic uremic syndrome (HUS) have been attributed to strains of serotype O157: H7. However, in some countries, the importance of non-O157 STEC as causes of HUS, HC, and other gastrointestinal disease is being increasingly recognized, particularly the serogroups O26, O103, O111, O121, O45, O145, and O91 **(Farrokh *et al.*, 2013).**

Illness caused by enterohaemorrhagic ***E. coli*** can range from self-limited watery diarrhea to life-threatening manifestations such as hemorrhagic colitis, hemolytic uremic syndrome, thrombocytopenic purpura and may lead to death **(alexander and Prado, 2003)**.

Recently there is additional concern about the presence of *E. coli* in cheeses because of the ability of some strains to cause foodborne illness and lead to a high incidence of food outbreaks **(Chapman, 2000)**.

As milk is of high nutritive value containing vitamins, minerals, and proteins so it promotes the growth of ***E. coli*** and other many microorganisms as a result of contamination from using unsterilized utensils, improper preparation of dairy animals, uses of contaminated water supplies, or careless method of handling **(Trevena *et al.*, 1999)**.

Kareish cheese is considered one of the excellent sources of protein, phosphorus, calcium, and many trace elements and due to its high nutritive value and low prices, it is considered as a popular food in Egypt. It is

manufactured from skim milk naturally fermented by lactic acid bacteria at home, so it may contaminate from sources during is manufacture, handling , and distribution that renders it a source of infection with many diseases **(Thabit, 2003)**.

Therefore, evaluation of the hygienic quality of Kareish cheese manufactured in el Gharbia governorate if one of the objects of this study.

Although milk pasteurization removes almost all pathogenic *E. coli*, inactivation of ***E. coli*** Shiga toxins has not been proven. However, following the consumption of pasteurized milk, an outbreak in North Cumbria, England, during 1999 with HUS cases have been reported and no living bacteria were found in the milk samples **(Goh *et al.*, 2002).**

Epidemiological studies suggest that nonO157 Shiga toxin-producing ***E. coli*** (STEC) is a major player associated with foodborne disease outbreaks **(Johnson *et al*., 2006).**

The ten most clinically relevant STECs belong to serogroups O26, O103, O111, O145, O157, O91, O113, O128, O45, and O121 have been reported in milk and dairy products causing severe illness in humans including diarrhea, hemorrhagic colitis, and hemolytic uremic syndrome and may result in death (**Farrokh *et al*.,2003).**

Karish cheese serves as an ideal medium for STEC bacterial growth due to the synergistic effect of different factors that exist in cheese as (low pH, water activity, NaCl, and starter culture) **(Galvez *et al*., 2007)**.

In this work the detection of ***E. coli*** in soft cheeses was performed by isolation and identification of **Enteropathogenic *E. coli*** strains serotypes.

In the present study a total number of 150 samples of Karish cheese, Tallaga cheese, Domiatte cheese (50 samples of each) were collected from different localities in el – Gharbia governorate, Egypt, and examined for the

presence of *E-coli.* The Tallaga and Domiatte cheese from cheese dairy shops and Kareish cheese from both dairy shops and street vendors.

The incidence of ***E-coli*** in Kareisg cheese and Domiatte Cheese and Tallaga cheese proved by Selective culturing on (Eosin Methylene blue media) were isolated and identified as ***E –Coli*** as fellow: Table (3)

1- Kareish cheese:

Inspection of the table before revealed that *E. coli* could be isolated from examined Kariesh Cheese samples with an incidence rate of 54%. A nearly similar finding was reported by El-bajory *et al.*, (2015), El- nahas., *et al* (2015), Nagah *et al.*, (2012). a higher incidence was reported by Ibrahim *et al.,* (2019), Ombark *et al.*, (2016), El-kosi (2001), Mohammed (2015) while the lower incidence was reported by El-Sayed *et al.,* (2011), Zeinhom (2011), Hosney *et al.*, (2011), Hefner (2012), Abd EL- Latif, Ahmed (2012), Brooks, *et al.*, (2012), Nagah *et al.*, (2012), Samaha *et al.*, (2012), Elhadidy and Mohammed (2013), Bonyadian *et al.,* (2014), Abo Zeed (2014), Metwally and Fatma (2015), El Nahas *et al.*, (2015), Gamal *et al.*, (2 015),

Jehan *et al.,* (2015), Gamal and Soad (2016), Sameh (2016) and

Lotfy *et al.,* (2017).

2-Domiatti cheese:

It is evident from the table before that ***E. coli*** could be isolated from examined Damietta cheese samples with an incidence rate of 42%. A nearly similar finding was reported by **Ozen *et al.*, (2011), Ibrahim *et al.*, (2019) while the lower incidence was reported by Zeinhom (2011), Hefney (2012), Nagah *et al.*, (2012), Elhadidy and Mohammed (2013), Gamal and Soad (2016), Jehan *et al.*, (2015), Gamal *et al.*, (2015), Elbassary (2006), El-bajory ,*et al.*, (2015) and Lotfy *et al.*, (2017).**

3-Tallaga cheese:

It is evident from the table before that ***E. coli*** could be isolated from examined Tallaga cheeses samples with an incidence rate of 42%.

A nearly similar finding was reported by **El-Sayed *et al.*, (2011),** while alower incidence was reported by**, Hefney (2012),** Contamination of milk and milk products, with ***E. coli*** is largely due to unhygienic condition during production, processing, handling, and distribution. ***E. coli*** is a good indicator of fecal pollution as it exists in the normal microflora of the intestinal tract of humans and warm-blooded animals **(Olfa *et al*., 2013).**

2-Serological typing of isolated *E. coli* strains: Table (4)

It is evident from the results recorded in a table (5) that some of the isolated ***E. coli*** strains (64 strains) could be serologically typed into; Enteropathogenic (EPEC), Enterohemorrhagic (EHEC), Enteroinvasive (EAEC) included O119, O111, O86, O18, O128, O124**.**

The Serologically identified pathotypes of *E. coli isolates* from some types of soft cheese were:

(O111) **(EPEC)** Enteropathogenic ***Escherichia coli***, (O86) (**EHEC)** Enterohemorrhagic ***Escherichia col*i**, (O 18) (**Ex PEC)** extraintestinal pathogenic *Escherichia coli*, **(O128) (Atypical EPEC) Atypical EPEC** *Escherichia coli*, (**O124) (EIEC**) Entero Invasive ***Escherichia coli***. ***E. coli*** infections from contaminated cheeses surely constitute the publics hazards of living creatures especially in humans like meningoencephalitis, pyelonephritis, hepatitis, pneumonitis, pleurisy, pericarditis, peritonitis, osteomyelitis, appendicitis.

And even causes outbreaks and death of the consumers because cheeses eating without further heating for pasteurizations.

In this study, ***E .coli*** was isolated from 43% of samples, and 75% of isolates belonged to the EPEC serogroup. Recent studies in other countries indicate similar results. **Araujo *et al., (2002)*** detected high levels of fecal contamination in 95.5% of cheese samples in Brazil, and EPEC was isolated from 21.1% of the samples. In Egypt, 60 Karih cheese samples were examined **and *E. coli*** was isolated from 75% of the samples. Similarly, **Abbar and Kadder (1991**) reported that 40.5% of cheese samples in Iraq were contaminated with EPEC strains. In the present study, we detected 6 different EPEC serogroups in Egyptian soft cheese and that O127 was not found serogroup. Recent studies in other parts of the world indicate different prevalent EPEC serogroups in soft cheeses. In Brazil, O127 was the most frequently found serogroup, followed by O55 and O26 (**Araujo et al., 2002). Ahmed et al., (1988)** showed the most prevalent serogroups in Egypt are O111, O126, O128, O26, O25, and O125. In Iraq, four EPEC serogroups, including O111, O86, O125, and O119, are commonly isolated from cheese samples (**Abbar, 1988**). In this study, EPEC serogroups were detected as O119 serogroup. Similar to our results, **Abbar(1988)** reported that O119 is one of the prevalent EPEC serogroups in cheese. In Australia, O119 serogroup was isolated from 161 patients with diarrhea symptoms due to the consumption of soft cheese (**Crossly *et al.*, 1995).**

In another study in the west of Iran, ***E. coli*** was isolated from 51% of domestic Iranian soft cheese samples (**Shridhar *et al.*, 2004).** It appears that the rate of contamination of soft cheese with ***E. coli*** varies in different parts of Iran.

According to the results of this study, 19.48% of isolated ***E. coli*** strains from domestic soft cheese belong to EPEC strains.

3-Sensitivity test: Table (6)

The best antibiotics for the treatment of colibacillosis infections in animals and humans were: Ciprofloxacin, Enrofloxacin, Oxytetracycline respectively to ovoid the antimicrobial resistance. The presence of antimicrobial resistance with virulence genes among STEC strains is worrisome. Therefore, when the relationship between virulence genes and antimicrobial resistance occurs the antimicrobial resistance might be selected through the antimicrobial use, and as a result, the virulence genes would be selected as well.

It is evident from the table before that ***E. coli*** virulence genes of pathogenicity that causes public hazard on cheeses consumers even cause could be discussed as follows: Table (2,7) :

1- ***Stx1*** gene It is a Shiga toxin potent exocytotic and causes severe illness:

Its presence in our isolates by PCR was zero % and not found in all ***E.coli*** isolates Nearly similar findings were reported by **Taha *et al.*,(2019*)*** while higher incidences findings were reported by **Ibrahim *et al .,(2019),* Sabrey *et al.*,(2008), Hussien *et* al., (2019), El-hadidy and Mohamed (2012), Embarks *et al.,* (2016), Cholesteric *et* al.,(2017).**

2- ***Eae A*** gene it an intimin secretion gene that enhance attachment and colonization:

Its presence in our isolates by PCR was 28.5 % of ***E. coli*** isolates: while lower incidences findings were reported by **Ibrahim *et al., (2019),* Sabrey *et al.,***

***(2008),* Hussien *et al., (2019*), El-hadidy and Mohamed *(2012),* Chalechtori *et al.,* (2017).**

3- ***Fim H gene*** it is responsible for biofilm formation and enhance infection of Uropathogenic *Escherichia coli* and the structural rule of type 1 fimbriae and responsible of about 92 % of the pathogenicity and diseased cases in human patients suffering from Urinary tract infections: Its presence in our

isolates by

PCR was 100 % of *E. coli* isolates: similar results and findings were reported by **Garofalo *et al.*, *(2007)*** while lower incidences findings were reported by**, Watts *et al.*, *(2010),* Mladin *et al.*, (2009), Arabi *et al.*, (2012), Tarchouna *et al.*, (2013), Sabrey *et al.*, (2008), Zohreh *et al.*, (2014), Ibrahim *et al.*, (2019).**

the current study supports the finding that raw milk and various milk products can be regarded as critical sources of pathogenic *E. coli* This explains the need of strict monitoring and surveillance for effective measures of hygiene and sanitary practice during the production of milk and various milk products.

CONCLUSION AND RECOMMENDATIONS

6- Conclusion and Recommendations

Current Study Supports The Finding That Soft Cheese Can Be Regarded As Critical Source Of Pathogenic *E. coli.* This Explains The Need Of Strict Monitoring And Surveillance For Effective Measures Of Hygiene And Sanitary Practices During the Production Of Soft Cheeses (Kariesh, Domiatti, Tallaga). The obtained results in this work give pieces information that some of the examined soft cheese like Kareish Cheese, Damietta cheese ,and Tallaga Cheese Samples Sold in the Gharbia Governorate Were Of Poor Microbiological Quality And That Indicate Inadequate Hygienic Measures Adopted During Their Production, Handling, Processing, Packaging And Distribution Of The Examined Soft Cheeses Products.

Contamination Of The Examined Kareish Cheese, Damietta cheese ,and Tallaga Cheese Samples With Pathogenic Microorganism with *E.coli* is a reliable index of fecal pollution, poor personal hygiene, constitute a public health hazard can range from self-limited watery diarrhea to life-threatening manifestations such Hemorrhagic Colitis, Hemolytic Uremic Syndrome, Thrombocytopenic Purpura, Meningoencephalitis and may lead to death and may lead to food poisoning and undesirable changes of the products that render them unfit for human consumption that resulted in Health And Economic Losses.

Therefore, to SafeGuard Consumers From Being Infected Preventing The Sailing Of This Types Of Soft Cheeses From Excellent Surveillances On The Land Of The Targeted Zones In The West Territories Otherwise Produced This Types Of Cheeses Under The Veterinary Supervisions And Applied The Good Hygienic Practices During The Cheeses Manufacturing Stages After Ensuring That The Milk Of No Microbial Loads.

Stipulated the cheeses packages must have all the details of manufacturers and certificates of hygienic practices certificates and must be of the current year.

In Conclusion, It Deems Necessary That Concerned Authorities Should Impose Regulations And Bacteriological Standards For Control Of Soft Cheeses Production And Handling Not Only For Detection Of Errors And Defects But To Ensures That The Errors Are Corrected And Defects Are Not Repeated And Ensure A maximum Of Safety To The Consumers.

SUMMARY

7- Summary

This study was carried out to investigate isolations and serological identification of pathological pathotypes of *Escherichia coli* from Kariesh, Tillage, and Damietta types of cheeses in El _Gharbia governorate; as well as molecular identifications of some virulence genes from the isolated strains. One hundred and fifty samples of soft cheese, including 50 of each Kareish cheese, Damietta cheese, and Tallga cheese were collected from supermarkets, dairy shops, and street vendors at El- Gharbia governorate.

Microbiological examination of the examined samples revealed that *E. coli* could be detected in 54%, 42%, and 32% of examined Kareish cheese, Damietta cheese , and Tallage cheese samples, respectively. Serological typing of isolated strains of *E. coli* proved that they belong to Six Serotypes : O1119, O111, O86, O18, O128, and O124).

The Serologically Identified Pathotypes Of *E. coli Isolates* From Some Types Of Soft Cheese Were: (O119) **(EPEC)** Enteropathogenic *Escherichia coli*, (111)

(EHEC) Enterohemorrhagic *Escherichia coli*,(O86:) Enteropathogenic *Escherichia coli*, (O 18) (**Ex PEC)** Extraintestinal pathogenic *Escherichia coli*,**(O128) (Atypical EPEC) Atypical EPEC** *Escherichia coli*, (**O124) (EIEC**) Entero Invasive Escherichia coli. *E. coli* infections from contaminated cheeses surely constitute the publics hazards of living creatures especially in humans likes meningeal encephalitis, pyelonephritis, hepatitis, pneumonitis, pleurisy, pericarditis, peritonitis, osteomyelitis an appendicitis.

And even causes outbreaks and death of the consumers because cheeses eating without further heating for pasteurizations.

On the other hand, molecular identifications of some virulence genes from the isolated strains. *stx1* gene was negative in all examind sample ; e*ae A* gene were detected in 28.5% of examind sample, while *fim H* gene was discovered in all samples.

In conclusion, isolation of EPEC serogroups from domestic soft cheese represents a potential, as well as an indication, of the presence of other enteropathogenesis. Although recent studies on virulence factors indicate that not all EPEC strains can attaching /effacing lesion, it is however believed that a high prevalence of contamination with EPEC strains increases the risk of infection for children, due to the consumption of domestic soft cheese. It seems that further epidemiological investigations are needed to reveal the importance of contamination in domestic soft cheese in this area of Egypt.

REFERENCES

8- References

Abd-Ehamid, A. M. (2012): Production of Functional Kariesh Cheese by Microencapsulation of Bifidobacterium adolescents ATCC 15704. Adv. J. Food Sci. Technol. 4 (2): 112-117, 2012.

Abou-Donia, S.A. (2008): Origin, history , and manufacturing process of Egyptian dairy products: An overview. Alex. J. Food Sci. Technol. 5: 51-62.

Ahmed, M. S. (2002): Polyamine pattern of some microorganisms isolated from environment milk and soft cheese. ph. D. Thesis, Fac. Vet. Med., Assuit Univ., Egypt, P. 55-64.

Alice, G.M G.; Ana, C.P. R; Joao, R.C. A. and Anita, T.(2000):

Enter pathogenicity markers in *Escherichia coli* strains isolated from soft white cheese and poultry in Rio de Janeiro, Brazil. Food Microbiology, 17(3): 321–328.

AL-Jeddah, J. H. and Robinson, R. K. (2001): The chemical composition and microbiological quality of some white brined cheeses on sale in Qatar.

Egyptian J. Dairy Sci., 29: 325.

Allen, K. J; Lapp, D; McKellar, R.C; Griffiths, M.W. (2008): Examination of stress and virulence gene expression in EscherichiacoliO157:H7 using targeted microarray analysis. Foodborne Pathos Dis. (5): 437–447.

Almeida, G.; Figueredo, A.; Role, M.; Barros, R.M.; Gibbs, P.; Hogg, T. and Teixeira, P.(2007): Microbiological characterization of randomly selected Portuguese raw milk cheeses concerning with reference to food safety. J. Food Prot., 70(7): 1710-1716.

APHA American Public Health Association (2004): Where, H.M. and Frank, J.F. (eds.). Standard methods for the microbiological examination of dairy products, 17th Ed., Washington, D.C., USA.

Assupol, G.L.H; Cardozo, M.V; Berardi, L.G; Malyuta, R.P; Silva, J.T; Avila, F.A.D; McIntosh, D; Robalo, E.C.(2015): Antimicrobials resistance patterns and the presence of stx1, stx2 and eae A in Escherichia coli. Rev Bras Saudi Prod Anima (16): 308-16.

Aseel A. Saeed. (2015): Detection of E. coli isolated from cheese by virulence factors by PCR technique in Al- Diwaniyah city. Al-Qadisiya Journal of Vet. MedChi. Vol.15 No. (1): 134-138.

Arabi S, T. F; Nader S, N. A; Jafarpour, M, Naghshbandi, R. (2012): The common fimbriae genotyping in Uropathogenic Escherichia coli. *Ann Biol Res.* 3(10): 4951–4. Araujo,

Araujo, V. S., V. A. Pagliarea, M. L. Queiroz, A. C. Freitas-Almeida. (2002): Occurrence of *Staphylococcus* and enteropathogens in soft cheese commercialized in the city of Rio de Janeiro, Brazil. J. Appl. Microbiol. 92, 1172-1177

Bach, S. J; McAllister, T. A; Vieira, D. M; Gannon, V. P. J. , and Holley, R. A. (2002): Effect of monensin on survival and growth of Escherichia coli O157: H7 in vitro. Can. Vet. J. (43:718-719).

Banatvala, N., P. M; Griffin, K. D; Greene, T. J; Barrett, W. F; Bibb, J. H; Green,and J. G.Wells. (2001): The United States National Prospective Hemolytic Uremic Syndrome Study: Microbiologic, serologic, clinical, and epidemiologic findings. J. Infect. Dis**. (**183) :1063–1070.

ibaraheem, O. H.; Hoda, A. E.; Wafaa, M. B. and Naglaa, F. G. (2007): Bacteriological quality of some dairy products (karish cheese and ice cream) in Alexandria. Microbiology Department, High Institute of Public

Health, Alexandria University, Egypt. J. Egypt Public Health Assoc., 82(65) :492-510.

But, L; Miko, A; Krause, G; Pries, K; Haby, S; Steege, K; Albrecht, N. (2007): Identification of human-pathogenic strains of Shiga toxin-producing *Escherichia coli* from food by a combination of serotyping and molecular typing of Shiga toxin genes. Appl Environ Microbiol (73): 4769-

75.

Blanco, M., Blanco, J.E;Mora, A; Rey, J;Alonso, J.M;Hermoso, M; Hermoso, J;Alonso, M.P; Dhabi, G; Gonzalez, E.A;Bernardez, M.I.

(2003): Serotypes, virulence genes, and intimin types of Shiga toxin (verotoxin)- producing E. coli isolates from healthy sheep in Spain. J Clin Micro. 41(4):1351-1356.

Blass, I; Kholoud, M; Ismail, M. M. (2003): Effect of draining method on the quality of Kareish Cheese. Journal of Agriculture Science, Mansoura Univ. 28 (10): 7365-7374.

Bisi-Johnson, M.A.; Obi, C.L.; Vasaikar, S.D.; Baba, K.A. and Hattori, T. (2011): Molecular basis of virulence in clinical isolates of Escherichia coli and Salmonella species from a tertiary hospital in the Eastern Cape, South Africa. Gut Pathogens 2011, 3:9.

Bolton, D. J. (2011). Verocytotoxigenic (Shiga Toxin-producing) *Escherichia coli:* virulence factors and pathogenicity in the farm to fork paradigm.

Foodborne Pathogens and Disease, 8(3), 357-365. http://dx.doi.org/10.1089/fpd.2010.0699. PMid:21114423.

Bonyadian, M.; Moshtaghi, H.; Mariam, A. T. (2014): Molecular characterization and antibiotic resistance of enterotoxigenic and

enteroaggregative *Escherichia coli* isolated from raw milk and unpasteurized cheeses. Veterinary Research Forum, 5 (1): 29 - 34.

Brown, R. C. 2004. The Complete Book of Cheese, Chapter (3). Roquefort cheese dressing, bottled USA**.**

Brett, K.N; Hornitzky, M.A; Bettelheim, K.A; Walker, M.J; Djordjevic, S.P. (2003): Bovine non-O157 Shiga toxin 2-containing Escherichia coli isolates commonly possess stx2-EDL933 and/or stx2vhb subtypes. J Clin Microbiol (41) :2716-22.

Brooks, J.C.; Martinez, B.; Stratton, J.; Balanchine, A.; Kostroma, R , and Hutkins, R. (2012): Survey of raw milk cheeses for microbiological quality and prevalence of foodborne pathogens. Food Microbiol., 31(2):154-158.

Campos, L. C; Whitham, T.S; Gomes, T.A.T; Arnd, C.R.J; Trybulski, R.I. (1994): *Escherichia coli* serogroup O111 includes several clones of diarrheagenic strains with different virulence properties. Infect. Immun. (62): 3282-3288.

Center for Disease Control and Prevention. (2006). Importance of culture confirmation of Shiga toxin-producing *Escherichia coli* infection as illustrated by outbreaks of gastroenteritis - New York and North Caroline, 2005.

Morbidity and Mortality Weekly Report, 55(38), 1042-1045.

Cergole-Novella, M.C., Pignatari, A.C., Castanheira, M. and Guth, B.E. (2011) Molecular typing of antimicrobial-resistant Shiga-toxin-producing *Escherichia coli* strains (STEC) in Brazil. *Res. Microbiol*., 162(2): 117-123.

Claeys, W.L., Cardoen, S., Daube, G., De Block, J., Dewettinck, K., Dierick, K. and Herman, L. (2013) Raw or heated cow milk consumption: Review of risks and benefits. *Food Contr*., (31) : 251-262.

CLSI, Clinical and Laboratory Standards Institute. (2018) Performance Standards for Antimicrobial Susceptibility Testing; Twenty-Fourth Informational Supplement. Available from: http://www.microbiolabbg.com/wp-content/uploads/2015/05/CLSI-2014.pdf. Accessed on 20-032017.

Crossly, V. M. M; SimpsonI, M. F . (1995): *Echerichia coli* strains associated with gastroenteritis in infants with special reference to a hospital outbreak due to *E. coli* type O111, B4. Can. J. Microbiol. 1, 257-283.

DlsvikL, Q. Y. A. L; WASTESON, E.H. (1991): Pathogenic *Escherichia coli* found in food. Int. J. Food Microbiol. 12, 103-119.

Durant, C.V.D; Splitter (1992): Compendium of Methods for the Microbiological Examination of Food. 3^{ed} ed., American Public Health Association, pp. 325-367, 849-852.

Dias, M. A; Santana, A. Scrub, A. Glair, J. A. F; Fernandes, de Dliveira, C. Abuna, E. (2012): Dn the implementation of good manufacturing practices in a small processing unity of mozzarella cheese in Brazil. *Food Control*, 24(1-2), 199-205. http://dx.doi. org/10.1016/j.foodcont.2011.09.028.

Dipinto, L; Santa niello, A; Fontanilla, M; Lagos, K; Floretta, A; Mena, L.F. (2006): Presence of Shiga toxin-producing *Escherichia coli* O157: H7 in living layer hens. Lett. Appl. Microbiol. 2006;43: 293–295.

Duello, T; Delannoy, S; Gannet, S; Foch, P; Loukoumis, E; Montel,

M.C. and Sergentet-Thevenot, D. (2017): Molecular characterization of O157: H7, O26: H11 and O103: H2 Shiga toxin-producing *Escherichia coli* isolated from dairy products. *Int. J. Food Microbiol*., 253: 59-65.

Dong, T. and Schell horn, H.E. (2009): Global effect of Repos on gene expression in pathogenic Escherichia coli O157: H7strain EDL933.BMC Genomics10, 349.

Johnson K.E., Thorpe C.M. & Sears C.L. (2006): The emerging clinical importance of non-O157 Shiga -toxin producing Escherichia coli. Clin. Infect. Dis. (43): 1587-1595.

Dkura, M. H., & Marin, J. M. (2014). Survey of Minas fresco cheese from Southwest Minas Gerais for virulence factors and antimicrobial resistance in *Escherichia coli* isolates. *Ciência Rural*, 44(8):1506-1511. http://dx.doi.org/10.1590/0103-8478cr20131237.

Ebrahim, R; Cephems and Pouya, P. (2011): Prevalence and antimicrobial resistance of Escherichia coli O157 isolated from traditional cheese, ice cream , and yogurt in Iran**.** Afr. Microion. vol.5 (22) : 3706 - 3710.

El-Baradari, G.; Bechet, D. A. , and Ogier, J. C. (2007): Biodiversity of Bacterial Ecosystems in Traditional Egyptian Domiatti Cheese. J. Appl.

Environ. Microbiol., 73(4):1248- 1255.

El-Kosi, O. H. R. (2001): Occurrence of some enteric pathogens and their indicators in some Egyptian raw milk products. Assiut Vet. Med. J. 45(89): 48-74.

El Sayed, M.A.; Hosny, I.M.; El Kholy, W.I.; El Dairouty, R.K. and Sahar, H. S. M. (2011): Microbiological evaluation of Egyptian white soft cheeses style. Journal of American Science, 7(5):517-526.

Elba gory, A. M; Eman, Sh. E and Eman, K.F. (2015): Impact of Probiotic Strains on Growth of Some Food Poisoning Bacteria from Milk and Soft Cheese.NutrFoodTechnol1(2): doihttp://dx.doi.org/10.16966/247060

86. 107.

El-Bassey, M. M. A. (2006): Sanitary status of milk and some milk products marketed in suburbs of Assiut Governorate. Ph.D. Thesis, Fac. Vet. Med., Assiut Univ.

Elhadidy, Mohammed, M.A. (2013): Shiga toxin-producing *Escherichia coli* from raw milk cheese in Egypt: Prevalence, molecular characterization, and survival to stress conditions. Lett Appl Microbiol (56): 120-7.

El Hofi, M; Ismail, A; Abd Rabo, F; El-Dieb, S; Ibrahim, O. (2010): Studies on the acceleration of Ras cheese ripening by aminopeptidase enzyme from Delvalle's pancreas utilization of buffaloes' pancreas aminopeptidase in an acceleration of Ras cheese ripening. New York Sci J (3): 91-6.

EL Kabel, M. A. I; Hend, A. E; Nahla, A. S and Islam, A. E. (2020): Survival of Shiga-toxin Producing *Escherichia coli* (STEC) during the storage of Karish cheese. AJVS. Vol. 64 (1): 47-53.

El nanas, A. W1.; Mohamed, H. A; El barbary, H. A2. And Mohamed H. S1. (2015): Incidence of E. coli in raw milk and its products. Bennah Veterinary Medical Journal, Vol. (29), NO. 1:112-117.

EL-holy, M; EL-sinewy, S.H; Meshref, A.M.S and Korany, A.M

(2014) : Microbiological Quality Of Domiatti Cheese And The Influence Of Probiotics On The Behavior Of Staphylococcus Aureus And Escherichia Coli O 157: H 7 In Domiatti Cheese. Journal of Food Safety. (34): 396–406.

Esraa Gaber Hefney Ahmed (2012): Microbiological Evaluation Of Soft Cheese Sold In Beni – Surf Markets. For. PD. Degree.

Espié, E; Vaillant, V; Mariani-Kurkdjian, P; Grimont, F; Martin-Schaller, R; De Valk, H et al. (2006): Escherichia *coli O157* outbreak associated with fresh unpasteurized goats' cheese. Epidemiology and Infection, (134) : 143–146.

Ekbal, M.A. I; Hend, A. E; Nahla, A. S and Islam, A. E. (2109): Incidence and molecular characterization of *Escherichia coli* in some dairy products. BVMJ 37 (1): 102-106.

Fadel, H.M. and Jehan, I. (2009): Prevalence and significance of *Staphylococcus aureus* and Enterobacteriaceae species in selected dairy products and handlers. Int. J. dairy Sci., *(*4): 100-108.

Farhad, S.C ; Navid, M.A; Esmat, A; Afshin, N. and Reza, S.C. (2017): Molecular characterization of Escherichia coli recovered from traditional milk products in Kashan, Iran. J. Vet World., (10): 1264 – 1268.

Farrokh, C., Jordan, K., Auvray, F., Glass, K., Dppegaard, H., Raynaud, S., Thevenot, D., Condron, R., De Reu, K., Govaris, A., Heggum, K.,

Heyndrickx, M., Hummerjohann, J., Lindsay, D., Miszczycha, S., Moussiegt, S., Verstraete, K., & Cerf, D. (2013): Review of Shigatoxinproducing *Escherichia coli* (STEC) and their significance in dairy production. *International Journal* Of Food Sci. Technol, Campinas, 37(1): 41-44, Jan.-Mar. 2017

Farrokh, C., Jordan, K., Auvray, F., Glass, K., Oppegaard, H., Raynaud, S., Thevenot, D., Condron, R., De Reu, K., Govaris, A., Heggum, K., Heyndrickx, M., Hummerjohann, J., Lindsay, D., Miszczycha, S., Moussiegt, S., Verstraete, K., and Cerf, O. (2013): Review of Shiga-

toxin producing Escherichia coli (STEC) and their significance in dairy production.

Int. J. Food Microbiol. 162(2) 190–212.

Francois, Z. N; Ahmed, N; Felicity, M. T; El-Soda, M. (2004): Effect of ropy and capsular exopolysaccharides producing strain of Lactobacillus Plantarum 162 M on characteristics and functionality of fermented milk and soft Kareish type cheese. Afr. J. Biotechnology. (3): 512-518.

Friedman, M; Shenika, P.R; Mandrel, R.E. (2002): Bactericidal activities of plant essential oils and some of their isolated constituents against *Campylobacter jejune*, *Escherichia coli*, *Listeria monocytogenes*, and *Salmonella Interic*. J Food Protect (65):1545-60.

Galvez, A; Gabriele, H; Lopez, R.L; Ben Omar, N. (2007): Bacteriocin based strategies for food biopreservation. I. J. of Food Microbiol. (120): 51-70.

Johnson, K.E; Thorpe, C.M; Sears C.L. (2006): The emerging clinical importance of non-O157 Shiga -toxin producing Escherichia coli. Clin. Infect. Dis. (43): 1587-1595.

Garofalo, C.K, Hooton, T.M; Martin, S.M; Stamm, W.E, Palermo, J.J; Gordon, J.I.(2007): Escherichia coli from the urine of female patients with urinary tract infections is competent for intracellular bacterial community formation.

Infect Immun. 75(1): 52–60**.**

Garbaj, A.M; Awad, E.M; Azwai, S.M; Abolghait, S.K; Naas, H.T; Moawad, A.A; Gammoudi, F.T; Barbieri, I; Eldaghayes, I.M. (2016): Enterohemorrhagic *Escherichia coli* O157 in milk and dairy products from Libya: Isolation and molecular identification by partial sequencing of 16S rDNA. Vet World (9):1184-9

Goh, S.; Newman, C.; Knowles, M.; Bolton, F.J.; Hollyoak, V.; Richards, S.; Daley, P.; Counter, D.; Smith, H.R. and Keppie, N. (2002): E. coli O157 phage type 21/28 outbreak in North Cumbria associated with pasteurized milk.

Epidemiol Infect. 129(3): 451-457.

Ghanbarpour and Salehi (2010): Determination of Adhesin Encoding Genes in *Escherichia coli* Isolates from Omphalitis of Chicks. American Journal of Animal and Veterinary Sciences 5 (2): 91-96, 2010.

Gundogan, N. and Avci, E. (2014): Occurrence and antibiotic resistance of *Escherichia coli*, *Staphylococcus aureus*, and *Bacillus cereus* in raw milk and dairy products in Turkey. International Journal of Dairy Technology, 67 (4): 562- 569. 47

Gamal. A. I.; Osama, M. S and Azzat B.A. (2015): Microbiological Quality of Commercial Raw Milk, Domiati Cheese , and Kareish Cheese. Middle East Journal of Applied Sciences. Volume: (05) Pages: 171-176.

Gamal, M. H. , and Soad, M. G. (2016): Microbiological Quality of Soft Cheese Marketed in Cairo and Giza Governorates. Alexandria Journal of Veterinary Sciences. 50 (1): 18-23.

Gómez, D.; Miliwebsky, E.; Fernández. P. C; Baschkier, A.; Manfredi. E.; Zotta, M.; Nario, F ; Piquín, A.; Sanz, M.; Etcheverría, A.; Padola, N.; Parma, A.; and Rivas, M. (2002): Isolation and characterization of Shiga-toxin-producing Escherichia coli from frozen hamburgers and soft cheeses. Rev Argent Microbiol.; 34(2):66-71.

Hu, Q.T.u; Han, J; Zhu, X; Ding, Y; C,Yu, C. S. (2011): Development of multiplex PCR assay for rapid detection of Riemer Ella antipasti, *Escherichia coli*, and *Salmonella enteric* simultaneously from ducks. J. Microbiol. Methods, (87): 64–69.

Hamad, M.N.F. (2015): Effect of adding Gluco-Lactone, Different of Starter, Rennet on the chemical composition, Yield and economic study of Kariesh cheese. Int.J. Food. Sci. Nut. Eng. (5) : 130 – 40 .

Hegazy, N.M; Nasr, M.M; Fayed, A.E; Youssef, M.S. (2012): Economics scale for processing of white soft cheese in Egypt. Egypt. J. Agric. Econ. (22): 1079-1094.

Honish, L., Predy, G., Hislop, N., Chui, L., Kowalewska-Grochowska, K.,

Trottier, L., et al. (2005): An outbreak of E. coli O157: H7 hemorrhagic colitis associated with unpasteurized gouda cheese. Canadian Journal of Public Health, 96, 182–184.

Hosny, I.M.; El Kholy, W.I.; Murad, H.A. and El Dariot, R.K. (2011): Antimicrobial activity of Curcumin upon pathogenic microorganisms during manufacture and storage of a novel style cheese 'Karishma. Journal of American Science, 7(5):611-618.

Hussain, A.; Ali, Sh.; Shakira, H.A.; Qazia, J.I.; Alic, Kh. and Ullahb, N. (2014): Prevalence of *Escherichia coli* in milk and different milk products sold under market conditions at Lahore. Scientific Journal of Animal Science, 3(1): 22-26.

Heba, H; Ayman, E; Marwa, S; Ghada, H; Ihab, M; Turki, D; Ayman, M and Eman, M. (2019): Molecular characterization of Escherichia coli isolated from cheese and biocontrol of Shiga toxigenic E. coli with essential oils.Ital.j.Food.Saf. 8 (3) : 8291.

ICMSF "International Commission on Microbiological Specification" (2006): Microorganism in Foods, Microbial ecology of food commodities. 2nd ed. Kluwer Academics, Plenum Publishers. Lender's, U. K.

Jafari, A; Alani, M; Bouzari, S. (2012): *Escherichia coli*: A brief review of diarrheagenic pathotypes and their role in diarrheal diseases in Iran. Int J Manga (4): 102-17.

Jehan, I. I. ; Ehab , S. ; Ahmed, S. and Alaa, A. H. (2015): Microbial Quality of Some Dairy Products in Ismailia City. 2nd Conference of Food Safety, Suez Canal University, Faculty of Veterinary Medicine Volume(I), Page 14-21.

Johnson, K.E; Thorpe, C.M; Sears, C.L. (2006): The emerging clinical importance of non-O157 Shiga- toxin -producing Escherichia coli. Clin. Infect. Dis. (43): 15871595.

Keene, W. E. K; Hedberg, D. E; Herriot, D. D; Hancock, R. W; McKay, T. J; Barrett, and Fleming, W.D. (1997): A prolonged outbreak of *Escherichia coli* O157: H7 infections caused by commercially distributed raw milk. J. Infect. Dis. (176):815–818.

Koeman, E. W., Allen, S. D., Schneckenburger, P., Janna, C., & Winn, I. W. C. (1997). *Color atlas and textbook of diagnostic microbiology* (5th ed.).

Philadelphia: Lippincott Company.

Karnak, J. Johnson, J.L. (2001): Enterobacteriaceae, *coliforms*, and *Escherichia coli* as quality and safety indicators. In: Downes, F.P., Ito, K. (Eds.), Compendium of Methods for the Microbiological Examination of Foods, fourth ed. American Public Health Association, pp. 69-82.

Karmali, M.A; Gannon, V; Sergeant, J.M. (2010): Verocytotoxin producing *Escherichia coli* (VTEC). Vet. Microbiol. (140): 360–370.

Kosta, M; Matras, M; Skandalis, P; Drowsing, E.H. (2010): Prevalence and sources of cheese contamination with pathogens at farm and processing levels. Food Control (21) : 805 – 15 .

Lara, V.M; Caremark, A.B; Santeria, D.F; de Sa, M.F; Santurio, J.M; Alves, S.H. (2016): Antimicrobial susceptibility of Escherichia coli strains isolated from Alouatta spy. feces to essential oils. Evid Based comply Altern Med 10: 30.

Ladan, M. N, and Reza, G. (2006): A study on enteropathogenic Escherichia coli isolated from domestic Iranian soft cheese. Veterinary ski Archiv 76 (6): 531-536.

Lopez-Exposito, I; Amigo, L; Recio, I. (2012): Amini- review on health and nutritional aspects of cheese with a focus on bioactive peptides. Dairy Sci Technol (92): 419- 38.

Lotfy, A. A.; EL-Touhy, M. E. and Al- Ashmawy, M. A. (2017): Assessment of raw milk and some dairy products for the presence of Escherichia coli with special reference to E. coli O157: H7 and their control using natural plants. Annals of Veterinary and Animal Science. vol 4(1).

Lothar,B ; Angelika, M; Gladys , K; Karin,P ; Sabine , H; Katja, S and Nadine, A. (2007): Identification of Human Pathogenic Strains of Shiga Toxin- Producing Escherichia Coli from Food by Combination of Serotyping and Molecular Typing Of Shiga Toxin Genes. app. Environ.Microbiol.73(15): 47694775.

Mac Fadin, J. F. 2000. Biochemical tests for identification of medical bacteria. 3rd Ed. Lippincott Williams and Wilkins, Washington.

Marier, R; Wells, J. G; Swanson, R. C; Callahan, Wand Mehlman, I. J. (1973): An outbreak of enteropathogenic *Escherichia coli* foodborne disease traced to imported French cheese.*Lancet*,2(7842),13761378.http://dx.doi.org/10.1 016/S0140-6736(73)93335-7.

Marvin, L. S. (1976): Compendium of Methods for the Microbiological Examination of Food. 2nd ed., American Public Health Association, pp. 115.

Metwally, A. M. M. and Fatma, H. M. Ali (2015): Escherichia coli in Dairy Products From Retailers And Small Dairy Shops. J. Food and Dairy Sci., Mansoura Univ., Vol. 6 (5): 349 – 355.

Mladin, C; Usein, C.R, Chifiriuc, M; Palade, A; Slave, C.L; Negut, M. (2009): Genetic analysis of virulence and pathogenicity features of uropathogenic Escherichia coli isolated from patients with neurogenic bladder. *Rom Biotech Lett.* 14(6): 4906–11.

Mohamed, S.H.S; Zaky, W.M; Kassem, J.M; Abbas, H.M; Salem, AL-AhI H.A.H; Abbas, H.M; Salem, M.M.E; AlAHAH.(2013): Impact of antimicrobi al properties of some essential oils on cheese yogurt quality. Wrld Appl Sci J (27): 497 – 507.

Mohamed. E and Mohamed .A.M. (2012): Shiga Toxin-Producing Escherichia Coli From Raw Milk Cheese in Egypt: Prevalence, Molecular

Characterization andSurvivaltoStressConditions.J.Applied.microbiology.Do 1:10.1111/Iam 2023.

Mohamed, E. A. S. (2015): Studies on enteric pathogens in milk and some dairy products with special reference to *Salmonella*. M.V.Sc. Thesis, Fac.

Vet. Med., Beni-suef Univ.

Moustafa, M. F. (2004): Studies on *E. coli* O_{157}: H_7 in milk and some milk products. Ph. D. Thesis, Fac. Vet. Med. Assiut Univ., Egypt.

Muniesa, M; Blanco, J.E; De Simon, M; Serra-Moreno, R;Blanch, A.R;

Jofre, J. (2004): Diversity of stx2 converting bacteriophages induced from Shiga-toxin producing Escherichia coli strains isolated from cattle.Microbiology (150): 2959–2971.

Nagachinta, S. and Chen, J. (2008): Transfer of class 1 integron-mediated antibiotic resistance genes from Shiga toxin-producing *Escherichia coli* to a susceptible *E. coli* K-12 strain in stormwater and bovine feces. *Appl.*

Environ. Microbiol., 74(16): 5063-5067.

Nagah, M. S.; Sabreen, M.S.; Wallaa, F. A. and Mira, K. G. (2012): Prevalence of *Escherichia* and other Escherichia species in raw milk and some dairy products in Assiut City, Egypt. Journal of American Science, 8(11):333-341.

Najand, L. M. and Ghanbarpour, R. (2006): A study on Enteropathogenic *Escherichia coli* isolated from domestic Iranian soft cheese. Veterinary ski Arhiv., 76 (6): 531-536.

Nabil, G; Franconieri, I ; Basanisi , G.M ; La Bella, Tozzoli, R; Caprioli, A and Lasalandra, G. (2016) : Isolation of shiga toxin-producing Escherichia coli in raw milk and mozzarella cheese in southern Italy . jornal of dairy science. Volume (99) : 7877-7880.

O'Brien, A. D., and J. B. Kaper. (1998): Shiga toxin-producing *Escherichia coli:* Yesterday, today, and tomorrow. Pages 1–11 in *Escherichia coli* O157: H7 and other Shiga toxin-producing *E. coli* strains.

J. B. Kaper and A. D. O'Brien, ed. ASM Press, Washington, DC.

Mubarak. R. A; Hinenoya, A; Awasthi, S. P; Iguchi. A; Shima, A; Elbagory, A. R; Yamasaki, S. (2016): Prevalence and pathogenic potential of Escherichia coli isolated from raw milk and raw milk cheese in Egypt. Int J Food Microbiol. 16;221 :69-76.

Orskov, F. and Orskov, I. (1973): Principles of Bacteriology, Virology , and Immunity. 7 edn., Vol.2, cited by Topley and Wilsons (eds)., (1984). Edward Arnold, pp. 288. 56

Oksuz, O.; Arici, M.; Kurultay, S. and Gumus, **T. (2004):** Incidence of *Escherichia coli* O157 in raw milk and white pickled cheese manufactured from raw milk in Turkey. Food Control, 15(6): 453–456.

Ortolani, M. B.; Yamazi, A. K.; Moraes, P. M.; Viçosa, G. N.and Nero, L. A. (2010): Microbiological quality and safety of raw milk and soft cheese and detection of autochthonous lactic acid bacteria with antagonistic activity against *Listeria monocytogenes*, Salmonella Spp., and *Staphylococcus aureus*. Foodborne Pathog. Dis., 7(2):175-180.

Ozen, K.; Several, S. K.; Erhan, K. and Ahmet, G. (2011): Microbiological quality of white pickled cheese produced in small plants in Burdur, Turkey. Journal of Food, Agriculture & Environment, 9(2): 110 - 112.

Paneto, B.R.; Schocken-Iturrino R.P.; Macedo, C.; Santo, J. M. E, and Marin, (2007): Occurrence of toxigenic Escherichia coli in raw milk cheese in Brazil. Arq. Bras. Med. Vet. Zootec., (59): 508-512.

Patercia, C and Jose, M.M.(2017): Occurrence of non-O157 Shiga toxin encoding *Escherichia coli* in artisanal mozzarella cheese in Brazil: risk factor associated with food workers.SCI.Technol(Campinas).37: No1

P**aton, J. C., & Paton, A. W. (1998).** Pathogenesis and diagnosis of Shiga toxin-producing *Escherichia coli* infection. *Clinical Microbiology Reviews*, 11(3), 450-479. PMid:9665978.

Pennington, H. (2010): Escherichia coli O157. Lancet, 376,1428–1435.PeterIkechukwu, J.O. A.I. 2014. Addition of Inulin and Manganese Chloride Increased Antibacterial Metabolites in Yoghurt. Pak. J. Nutr.(13): 12-16.

Pradel, N., Livrelli, V., De Champs, C., Palcoux, J. B., Reynaud, A., Scheutz, F., Sirot, J., Joly, B., & Forestier, C. (2000). Prevalence and characterization of Shiga toxin-producing *Escherichia coli* isolated from cattle, food, and children during a one-year prospective study in France. *Journal of Clinical Microbiology*, 38(3), 1023-1031. PMid:10698990.

Pizarro, M.A; Orozco, J.H; Degarbo, S.M; Calderon, A.E; Nardello, A.L; Laciar, A; Ruttler, M.E. (2014): Virulence profiles of Shiga-toxin-producing *Escherichia coli* and other potentially diarrheagenic *E. coli* of bovine origin, in Mendoza, Argentina. Braz. J. Microbiol. (44) : 1173–1180.

Pusz, P; Bok, E; Mazurek, J; Stosik, M; Chudzik, K. (2014): Type 1fimbriae in commensal *Escherichia coli* derived from healthy humans. Acta Biochem Polon (61) : 389-92.

Quinn, P.J; Carter, M.E; Markey, B.K; CARTER, R.G. (1994): Clinical Veterinary Microbiology. 1st ed., Wolfe Publishing, pp. 220-226.

Quinn, P.J; Carter, M.E; Markey, B.K; Donnelly, W.J.C. and Leonard, F.C. (2002): Veterinary Microbiology and Microbial diseases.

MPG.Book 1st ed. Bodmin, Cornwall, UK.

Q., Holien, J.K; Parker, M.W; Tauschek, M; Robins Browne, R.M (2017): Control of virulence gene expression by the master regulator, Cfa, in the prototypical enterotoxigenic *Escherichia coli* strain, H10407. Front Microbiol 8:1525.

Rey, J., Sanchez, S., Blanco, J.E., Hermoso de Mendoza, J., Hermoso de Mendoza, M., Garcia, A., Tejero, N., Rubio, R. and Alonso, J.M. (2006).

Prevalence, serotypes and virulence genes of Shiga toxin-producing *Escherichia coli* isolated from ovine and caprine milk and other dairy products in Spain. *Int. J. Food Microbiol.*, 107(2): 212-217.

Rattler, M.E; Renna, N.F; Balbi, L; Garcia, B; Guidone, L; Fernandez, R; Puig, O; Ortiz, A.(2006): Characterization of enteroaggregative *Escherichia coli* strains isolated from children with acute diarrhea, in Mendoza, Argentina. Rev. Argent. Microbiol. (34): 167–170.

Ryser, E. T. (2001): Public Health Concerns, Pages 397–545 in Applied Dairy Microbiology. E. H. Marth and J. L. Steele, ed. Marcel Dekker Inc., New York, NY.

Sambrook, J; Fritscgh, E, and Mentiates. (1989): Molecular cloning. A laboratory manual. Vol !, Cold Spring Harbor Laboratory Press, New York.

Said, M. R, and Fahmy, M. A. (1991): A survey of the incidence of *Staphylococcus aureus, Escherichia coli*, and *Bacillus cereus* in some types of Egyptian cheese in Assuit city. Assuit J. of Agric. Sci., 22(2):239- 246.

Schrade, J. P; Yager, J. (2001): Implication of milk and milk products in food disease in France and different industrialized countries. Int. J. Food Microbiol. (67): 1-17.

Silva, Z. N; Cunha, S.A; Lins, C.M; de. A. M. Carneiro, M.A.de; L. Almeida, I.A.C; Queiroz, L.M. (2001): Isolation and serological identification of enteropathogenic *Escherichia coli* in pasteurized milk in Brazil. Rev. Saude Publica.(35): 375-379.

Shidfar, R. F; AMERY, A; Sadeghifar, N. J. (2004): A survey on Aflatoxin M1 and Bacterial contamination on domestic Iranian cheese in Liam from 2002 to 2004. National Congress on Food Hygiene and Safety. 21-23 Dec. Yazd, Iran.

Sabry, A. H; Laila, M. E . (2008): Informally Raw Milk And Kariesh Cheese Investigation On The Occurrence Of Toxigenic Escherichia Coli In Qena City, Egypt With Emphasis On Molecular Characterization. Ass. Univ. Bull. Environ. Res. (11): 35-42.

Samaha, H. A.; Haggag, Y. N.; Nossair, M. A. and Mohammad, H. S. (2012): Using some recent techniques in the diagnosis of some zoonotic bacterial diseases transmitted through milk. Alexandria Journal of Veterinary Sciences, 35 (1): 11-21.

Salwa, S. T. (2003): Occurrence of Escherichia coli in milk and some milk products in Assuit City. M. V. Sc. Thesis, Fac. Vet. Med., Assiut Univ., Egypt. 75-82.

Sameh Awad. (2016): Microbial safety criteria and quality of traditional Egyptian Karish cheese. African Journal of Microbiology Research. Vol. 10(22): 804-812.

Shatha, T.A; Amina, N.J and Ayad, M.A.(2011): Isolation and **characterization** E. coli O157H7 from meat &cheese and detection of Stx1, Stx2, hlyA, eae A by using multiplex PCR. Miraz. The magazine for Biological Technologies. Volume 5. Second Edition: 15-24.

Shelaih, M. A. (1979): Microbiological studies on Egyptian soft cheese. Ph. D. Thesis, Fac. Vet. Med., Cairo Univ., Egypt, p. 81-91.

Spano, G; Goffredo, E; Beneduce, L; Tarantino, D; Dupuy, Aand Massa, S. (2003): Fate of *Escherichia coli* D157: H7 during the manufacture of Mozzarella cheese. *Letters in Applied Microbiology*, 36(2), 73-76.

Todd, E. C. D; Michaels, B. S; Smith, D; Greig, J. D; Bartleson, C. A. (2010): outbreaks where food workers have been implicated in the spread of foodborne disease. Part 9. Washing and drying of hands to reduce microbial contamination. *Journal of Food Protection*, 73(10): 1937-1955.

PMid:21067683.

Trachoma; Ferjani, A, Ben-Selma, W; Boukadida, J. (2013): Distribution of uropathogenic virulence genes in Escherichia coli isolated from patients with urinary tract infection. *Int J Infect Dis.*17(6): 450–3.

Vanaja, S.K: Springman, A.C: Besser, T.E; Whittam, T.S; Manning, S.D. (2010): Differential expression of virulence and stress fitness genes between Escherichia coliO157: H7strains with clinical or bovine-biased genotypes. Appl. Environ Microbiol. (76): 60–68.

Virar, P. K.; Nayak, J. B; Brahmbhatt, M. N and Thaker, H. C.(2013): Study on isolation, molecular detection of virulence gene and antibiotic sensitivity pattern of Escherichia coli isolated from milk and milk products.Doi: 10.5455/vetworld.2013.541-545.

Walid, M. E; Mona, A. Y; Salwa, F. A. (2015): **Molecular** characterization and stress tolerance of Escherichia coli isolated from dairy and dried milk-related products. International Journal of Food Science and Technology. (50): 136–142.

Wallace, J. S. (1999): The ecological cycle of *Escherichia coli* O157: H7. Pages 195–223 in *Escherichia coli* O157 in Farm Animals. C. S. Stewart and H. J. Flint, ed. CABI Publishing, New York, NY.

Wells, J. G; Shipman, I.D; K. D. Greene, K.D; E. G. Sowers, E.G; J. H. Green, J.H; Cameron, D.N; Downes, F.P; Martin, L.M; Griffin, P.M and O stuff, M.S. (1991): Isolation of *Escherichia coli* serotype O157: H7 and other Shiga-like-toxin-producing *E. coli* from dairy cattle. J. Clin. Microbiol. (29) :985–989.

Woteki, C. EandKineman, D.B. (2003): Challenges and approaches to reducing foodborne illness. Annu. Rev. Nutr. (23) :315–344.

Watts, R.E; Hancock, V; Ong, C.L; Vejborg, R.M; Mabbett, A.N; Totsika, M. (2010): Escherichia coli isolates causing asymptomatic bacteriuria in catheterized and non catheterized individuals to possess similar virulence properties. *J Clin Microbiol.* 48(7): 2449–58.

Xia, X; Meng, J; Mcdermott, P.F; Ayers, S; Blickenstaff, K; Tran, T.T; Abbot, J; Zheng, J; Zhao, S.(2010): Presence and characterization of Shiga toxin-producing *Escherichia coli* and other potentially diarrheagenic *E. coli* strain in retail meats. Appl. Environ. Microbiol.

(76): 1709–1717.

Zhao, T.M; Doyle, P; Shere, J and Garber, L. (1995): Prevalence of enterohemorrhagic *Escherichia coli* O157: H7 in a survey of dairy herds. Appl. Environ. Microbiol. (61) :1290–1293.

Zohreh, H; Behnam, Z; Morteza, H; Razieh, M; Abolfazl, G (2015): Detection of FimH Gene in Uropathogenic *Escherichia coli* Strains Isolated From Patients With Urinary Tract Infection Jundishapur J Microbiol. 2015 February; 8(2): e17520.

Ghanbarpour and Salehi (2010): Determination of Adhesin Encoding Genes in *Escherichia coli* Isolates from Omphalitis of Chicks. American Journal of Animal and Veterinary Sciences 5 (2): 91-96, 2010.

ARABIC SUMMARY

الملخص العربى

أجريت هذه الدراسة لمعرفة مدى تواجد ميكروب الإيشيريشيا القولونية وعزلها وتصنيفها من عينات الجبن القريش والدمياطى والثلاجة فى محافظة الغربية وكذلك التصنيف الجزيئى لبعض جينات الدراوة الموجودة بالعترات المعزولة حيث تم تجميع 150 عينة من الأجبان الطرية بمعدل 50 عينة من كل من الجبن القريش والجبن الدمياطى وجبن الثلاجة والتى تباع بالسوبر ماركت ومحلات الألبان والباعة الجائلين في محافظة الغربية.

أظهر الفحص الميكروبيولوجي للعينات اللاتي تم فحصهن أن ميكروب الإشريشيا القولونية قد تم اكتشافها في 54٪ و 42٪ و 32٪ من عينات الجبن القريش والجبن الدمياطى وجبن التلاجة التي تم
فحصهن على التوالي. وأثبت إختبار التصنيف السيرولوجى للعترات المعزولة من ميكروب الإشريشيا القولونية أنها تنتمي إلى ستة أنماط سيرولوجية هى :

O119، O86،O18،O128 ، O124 ،O111

وتم عمل إختبار حساسية على المزارع البكتيرية حيث وجد إن المضادات الحيوية وهى بالترتيب : السيبروفلوكساسين والإنروفلوكساسين والأوكسى تترسيكلين تؤثر وتثبط نمو المعزولات السابقة ومن ناحية أخرى ، تم عمل الدراسات الجزيئية لبعض الجينات من السلالات المعزولة (سبع معزولات) بإنزيم البلمرة المتسلسل كان الجين stxl سلبيًا في جميع عينات الفحص ؛ تم اكتشاف جين A eae في 28.5٪ من عينات الفحص ، بينما تم اكتشاف جين fim H في جميع العينات.

الملخص العربى

نبذة عن حياة الباحث

ولد الباحث فى التاسع من سبتمبر عام 1981 ميلاديا بمحافظة البحيرة بمصر

أنهى تعليمه الإبتدائى والإعدادى بمدرسة الدلجمون أبو بكر الصديق الإبتدائية و الإعداية بالدلجمون عام 1991 ميلاديا

أنهى دراسته الإعدادية والمتوسطة والثانوية بالمملكة العربية السعودية عام 1995 ميلاديا
أنهى دراسته الثانوية العامة فى مدرسة الدلجمون المشتركة بمحافظة الغربية بجمهورية مصر العربية عام 1997 ميلاديا

حصل على بكالوريوس العلوم الطبية البيطرية من كلية الطب البيطرى بجامعة طنطا فرع كفر الشيخ دفعةعام 2003 بتقدير عام جيدجدا وترتيبه السابع تراكمى

يعمل بمعهد بحوث الصحة الحيوانية بفرع طنطا

جامعة بنها
كلية الطب البيطرى بمشتهر
قسم البكتريا و المناعة و الفطريات

قرار لجنة الحكم والمناقشة

قررت لجنة الحكم والمناقشة المكونة من السادة الآتية اسماؤهم بعد جلستها اليوم الموافق ٢٠٢٠/٩/٢٧ منح ط.ب / **محمد محمود محمد يوسف وهيبه** درجة الماجستير في العلوم الطبية البيطرية تخصص البكتريولوجيا والمناعة والفطريات من كلية الطب البيطري بمشتهر جامعة بنها .

أعضاء لجنة الحكم والمناقشة :-

أ.د/ أحمد عفيفي عبد الغفار معروف	**رئيس خارجي**
رئيس بحوث معهد بحوث صحة الحيوان ببنها	.....
أ.د/ أشرف عواد عبدالتواب	**مشرف ومحكم**
استاذ ورئيس قسم البكتيريولوجيا والمناعة والفطريات	.....
أ.د / أمل مصطفي عيد	**مشرف ومحكم**
رئيس بحوث صحة أغذية معهد بحوث صحة الحيوان فرع طنطا	.....
أ.د.م / فاطمة إبراهيم الحوفي	**محكم**
أستاذ مساعد البكتريولوجيا والمناعة والفطريات	.....

كلية معتمدة 2013

جامعة بنها

كلية الطب البيطرى
قسم البكتيريا والمناعة والفطريات

تواجد ميكروبات التسمم الغذائى فى بعض أنواع الجبن الطرى فى السوق المحلى

رسالة مقدمة من

محمد محمود محمد يوسف وهيبه

بكالوريوس الطب البيطرى بكفرالشيخ جامعة طنطا عام 2003

لنيل درجة الماجيستير فى العلوم الطبية البيطرية

قسم البكتيريولوجى والمناعة والفطريات

تحت إشراف

الأستاذ الدكتور

أشرف عواد عبد التواب

أستاذ ورئيس قسم البكتيريا والمناعة والفطريات

كلية الطب البيطرى

جامعة بنها

الأستاذة الدكتورة

أمل مصطفى عيد

رئيس بحوث من مركز البحوث الزراعية بالقاهرة

معهدبحوث الصحة الحيوانية المصرى فرع طنطا

Subject: General Requests for a Hearing Sessions on Scientifics Facts and the Entitlements to Academics and Researchs Degrees

"I have made great efforts to organize scientific facts through voice-recorded lectures and seminars across laboratory scientific disciplines. These serve the pathways and empirical evidence that help elucidate the title of my experimental scientific theses, aiming to achieve the maximum level of knowledge and fruitfuls investigations to earn the highest standards of international scientifics merits. This is supported by local and international reviews from scientifically, legally, and globally recognized university scholars—leaving no rooms for doubts or suspicions in awarding experimentals academics degrees to those contributing to the advancements of Egyptians laboratory scientifics researchs.

Thank you for your patience, attentive listening, and prompt response."

"All scientific facts have been verified through electronic links and proofs of publications in local and international researchs via Digitals Objects Identifiers (DOIs). I sincerely thank you for All , your fairness in judgments and integrity in discussions."

Name of Assistant Researcher : Mohammed Mahmoud Mohammed Yousef Wehaba , ,Egypt State , Arabic race .

Mobiles contact details : whats app and telegram app :

+2 01287325739 , +2 01287325749 , +2 01040179247 .

Identity card number : 28109091800516 .

Passport number : A29252911
My driving license : 113622
My insurance number : 13375222
My registration inside veterinary syndicate:31679
My professional license inside veterinary syndicate : 31770
Working place:animal health research institute , agricultural research center , egyption government
My military number :2004578801966
My military weapons mobilization number: 590101003 . (Reserve first lieutenant)
My working code number: 8840
My working insurance number : 5100

A-Universal DOI Records (Zenodo)
:https://doi.org/10.5281/zenodo.15584047
https://doi.org/10.5281/zenodo.15583021
https://doi.org/10.5281/zenodo.19388186
https://doi.org/10.5281/zenodo.17866111
https://doi.org/10.5281/zenodo.17745368
https://doi.org/10.5281/zenodo.17579555

https://doi.org/10.5281/zenodo.17064403
https://doi.org/10.5281/zenodo.18252846

National Publication (Egypt – BVMJ)

Published in: Benha Veterinary Medical Journal (BVMJ)

DOI:

https://doi.org/10.21608/bvmj.2020.37011.1230.

EKB Article ID: 116440

Article Link:

https://journals.ekb.eg/article_116440.html.

https://1024terabox.com/s/1esaDa3yLAzVO6y_UanHEXA

https://docs.google.com/presentation/d/1GhzHKm94jslnJLCxh4mugaamu5Z1E2fX/edit?usp=drive_link&ouid=101122019937214507234&rtpof=true&sd=true

https://1024terabox.com/s/12E-zJ7MjXCQRmhsU0B-Wlw

https://mega.nz/file/0rADhADZ#et3jasNoftpi7titAarrvkv6wJXFe1GZwe4zeuv4G_M

https://fvtmbuedu-my.sharepoint.com/:p:/g/personal/m_waheba55689_fvtm_bu_edu_eg/IQBZ_tDa5zG2RImdv_70a2awATFPi_Yif0DWO4FglZAEBKY?e=3Dy2b5

m.waheba55686@fvtm.bu.edu.eg

ALL RIGHTS RESERVED .


K1
K2
K3
K5
D3